

MEDICAL AND SURGICAL REPORTER

No. 1765.

PHILADELPHIA, DECEMBER 27, 1890. VOL. LXIII.—No. 26.

CLINICAL LECTURES.

CLOSURE OF VULVA FOR VESICO-VAGINAL FISTULA.—FIBROID TUMORS.

BY WILLIAM GOODELL, M. D.,
PROFESSOR OF GYNECOLOGY, UNIVERSITY OF
PENNSYLVANIA.

Closure of Vulva.

Gentlemen: This woman had her first labor twenty-five years ago, and is now fifty years old. She lived then in the country, and was four miles from the nearest physician. She fell into labor—her first and only one—with a shoulder presentation. Much delay took place before the doctor reached her side, and then he found that the waters had escaped, and the shoulder was so jammed down that he could not perform version. He sent for help, and two brother physicians came to his aid; but they also found it impossible to turn the child and deliver the woman. By this time many hours had elapsed, and while they were debating the propriety of performing embry-ucia, the child was turned and delivered by the rare mechanism of spontaneous evolution. Of course the child was dead; but that was not the only misfortune; for the bladder and the surrounding pelvic tissues and organs had been so seriously compressed as to cause sloughing of the womb, the whole base of the bladder, the entire track of the vagina, and all of the urethra. The vagina nearly closed up by cicatricial contraction and the urine was constantly dribbling from her person, making her very raw, bad smelling and miserable. She consulted eighteen physicians; but nothing could be done for her. Finally she came to me in 1877; and I performed the following operation with her consent, telling her that I did not know whether it would succeed or not.

It consisted in making an opening between the vagina and the rectum—in other words, a recto-vaginal fistula. Then I closed up the whole vulvar opening, and thus converted the rectum into a bladder. At first the rectum so resented the presence of the urine, that she had to pass her urine every few minutes, and I was afraid that I would be compelled to reopen the vulva. But finally rectal tolerance took place, and she became able to hold her water in the rectum for six hours at a time.

For thirteen years she got on very well; but a few months ago, probably through something undigestible which she had swallowed—perhaps a bone—a small fistulous opening occurred near the arch of the pubic bone, from which the urine dribbles. So she has come to-day to have it closed up. She is an opium-eater, and she looked so pale that I had her urine examined. Much albumin was found; but whether it is secreted from her kidneys, or comes from the admixture of feces in her urine, I cannot say. But we shall take additional precautions while administering the ether; for ether is obnoxious to diseased kidney, especially when caused by compression from large tumors; and we may then have arrested secretion as a result. Under such a circumstance, namely, the presence of a large tumor, I usually give chloroform.

The operation to-day will consist in simply freshening the sides of the sinus, and then bringing the raw edges together. My assistant has just syringed out the fistulous track with a 1 : 2,000 sublimate solution, preparatory to the operation. But he forgot that the vagina communicates with the rectum, and that this antiseptic solution, while quite harmless in the vagina, is capable of doing much mischief in the rectum. So we wash out the rectum with large quantities of water, and thus get rid of the danger of having the poison absorbed.

This episode reminds me of another,

which filled me with great anxiety at the time, but at which I can now laugh heartily when I think of it. Early last summer I removed a malignant growth from the cervix uteri of a woman and ordered daily douches of a 1 : 2,000 solution of corrosive sublimate: that is to say, a tablet of seven and a half grains of corrosive sublimate was put into a quart of water which was injected by the nurse into the vagina. As the patient convalesced, she would sit on the commode and would herself insert the nozzle of the syringe into the vagina. But on one extremely hot evening she unconsciously introduced the nozzle into the rectum. The nurse, not knowing how very much astray the nozzle had gone, pumped away until very nearly the whole quart had been taken up. Suddenly the woman began to feel severe rectal pain, and then she realized what she had done. The nurse, although very greatly alarmed, was quite equal to the emergency. She sent in great haste for me, and at once injected quart after quart of water until the lower bowel was well washed out. I hurried the messenger to the nearest drug store for lime water, and then ran all the way to my patient's house, which was nearly half a mile distant. A quart of lime water was now thrown up into the bowel, as an additional precaution; but it was hardly necessary, as the previous repeated injections by the nurse had carried off the poison. Beyond a slight soreness of the rectum, which lasted for twenty-four hours, my patient suffered nothing. But what with the heat, the anxiety, and my hard run, my underclothing became so drenched with perspiration that I had to change them all when I reached home.

But, to return to our patient. It sometimes happens in these cases of violent injury to the soft parts from labor, that menstruation is permanently arrested. Exactly why this happens, I cannot say; but so it was with our patient when I operated on her in 1877; and I have repeatedly seen it in other cases. In one of these cases—that of an unmarried girl—I never could find the cervix uteri, so destroyed and matted together were all the pelvic organs from a neglected labor. Twice, in vain efforts to discover it by dissection, I got into the peritoneal cavity. As she had lost the whole base of her bladder, and also had a very large recto-vaginal fistula, caused by the general sloughing, I closed up her vulva, as in the patient before you. In her case also,

the rectum needed education to tolerate the presence of urine. But ultimately it made a very good bladder. If my memory is not at fault, these are the only two cases in which I was forced to close up the vulva. I have, however, on one occasion closed up the womb, or rather the cervix, in the bladder, so that the woman menstruated afterwards into the bladder and out through the urethra.

While talking, I have been denuding the edges of the fistula. These are coaptated by shotted wire sutures. In order to avoid all tension on the parts by the accumulation of urine in the vagina, I shall insert a short catheter into the rectum. These sutures will not be removed for at least a week; but if there should be a leakage from the wound they will be kept in for two weeks, so as to aid the granulating surface to close up. The wire used was No. 31, which is the finest used in gynecological work. My rule in the selection of wire is, to use the finest compatible with safety—the finest for plastic operations on the cervix and on the bladder; the heaviest for perineal tears.

The lesson which we get from this case is to act promptly in obstetric emergencies. My rule is always to interfere, and that promptly, in certain conditions. For instance, in a face presentation, if the occiput cannot be brought down to become the presenting part, I always turn. In a breech case, I make it the rule to bring down one leg; and I thus have the labor wholly under my control. In a shoulder presentation I always give ether, and turn as soon as possible, by the bimanual, or bipolar, method. The most common cause of vesico-vaginal fistula is delay in a head-first labor, after the engagement of the head. Therefore this lesion is more likely to happen in cases of moderate disproportion than in extreme ones; for in the latter the head cannot engage, and it has to be opened before it can delivered.

Medicinal Treatment of Fibroid Tumors.

I expected to operate before you upon this case; but, upon thoroughly examining this woman and learning her history, I have decided not to do so; I have changed my mind and come to this let-alone treatment for some reasons which I wish to impress upon you. The patient is an unmarried woman, forty-two years old, and was beginning to go as long as seven weeks without seeing her menstrual periods, when, no;

long ago she began to menstruate profusely every three weeks, or, at least, to have severe hemorrhages from the uterus at these intervals. A few weeks ago her physician discovered, and informed her that she had a tumor of the womb; and she, woman-like, wished to "have it taken out at once," and gave him no peace until he brought her to me. So by agreement I came over to see her yesterday and decided not to operate, or, at least, not to operate now, as I shall presently explain to you. She is forty-two years old; therefore, approaching the climacteric, and her monthly periods had begun to dodge, before these violent hemorrhages set in. Now what causes these floodings? Usually one of only two diseases: either a cancer of the cervix uteri, or a uterine fibroid. If she had been the mother of children, the probability would be that she had a cancer; because this disease usually comes from a cervical tear. But as our patient has never had a child, the tumor turned out to be a uterine fibroid, as under such circumstances it usually does. Old maids and sterile women are very liable to have fibroid tumors of the womb, and very unlikely to have a uterine cancer. Yes, I fully agree with the diagnosis of her physician. She has multiple fibroid tumors of the uterus. There are, at least, three of them that are sub-peritoneal and attached to the left side of the womb. By them the uterus has been pushed over to the right side, and the left ovary is probably much dislocated in relation to the surrounding parts. Now, if any operation were performed, it would have to be one of two, either of which would be difficult and dangerous. The more common of these two is oophorectomy, with the hope of stopping menstruation and thereby relieving the certainly periodical, and more or less constant congestion of the womb. But this operation would probably be difficult, if not impossible, on account of the size of these multiple tumors. For, sometimes the ovary in these cases has no pedicle, as it is drawn down in close relation to the tumor. Sometimes it is stretched to a mere ribbon or a mere cord several inches in length. Sometimes, indeed, it is so imbedded in the tumor as not to be enucleated safely. The other operation would be the removal of the whole mass—womb, tumors and appendages. But there is some risk attending this operation—a risk to life which her condition does not warrant. Moreover she does not live by

manual labor, and therefore she can take care of herself until the change of life takes place, which will bring about the cure of her disease. Nature cures, by causing atrophy and absorption of these growths, whose nutrition is interfered with by the cessation of menstruation.

To tide the woman over to that time, needs some treatment. As she has plenty of fat and blood, she is to have her bowels kept open by saline cathartics; such as Rochelle salts, Seidlitz powders and citrate of magnesia. Then she is to take after each meal twenty drops of the fluid extract of ergot and ten grains of ammonium chloride. The effect of the ergot will be to contract the muscular fibres of the uterus, and thus, by compressing the vessels, to lessen the blood supply of the tumor. Sometimes the womb contracts so firmly as to squeeze the tumor out of its uterine nest into the uterine cavity and expel it per vaginam. The ergot may cause headache, vertigo and nausea, through vaso-motor disturbance; and then the dose must be reduced, but not discontinued wholly. The ammonium chloride acts as a resorbent, and clinches the action of the ergot. For continuous treatment, it should not be given in tablet form; as it is likely then to settle itself in one portion of the stomach and dissolve there, irritating the mucous coat of that portion. Give it largely diluted in water, and then it will mix with the contents of the stomach and get still further attenuated. This treatment by ergot and ammonium chloride should be continued for months. I have seen great results follow the continued use of these medicines. The hypodermic administration of ergotine is still more efficacious; but it gives so much pain, so discolors the skin and is so liable to be followed by abscesses or very painful indurations, that I now rarely resort to it.

HOMEOPATHY IN RUSSIA.—Homeopathy is said to be spreading in Russia, especially in the upper social strata. Societies for the propagation of the Hahnemannian doctrines have recently been established at Tschernigow, Odessa and Warsaw. As has been noticed in other countries, the clergy are conspicuous among the supporters of the great medical heresy, and in Russia the military mind seems also to have an elective affinity for globules and infinitesimal dilutions. At Tschernigow, one of the founders of the new Society is a Bishop.

**HYDATID CYST OF THE LIVER.—
PLEURISY ASSOCIATED WITH
TUBERCULOSIS.¹**

BY JOHN H. MUSSER, M. D.,

ASSISTANT PROFESSOR OF CLINICAL MEDICINE IN THE
UNIVERSITY OF PENNSYLVANIA, PHYSICIAN TO
THE PHILADELPHIA HOSPITAL, ETC.

Obstructive Jaundice Due to a Hydatid Cyst.

Gentlemen: A little sooner than I anticipated I am able to demonstrate the morbid anatomy of the case of obstructive jaundice I brought before you last week. You no doubt remember the patient, whose previous history we did not know, except that his jaundice was of a moderately chronic duration, lasting for two months. The fact that he was jaundiced because of obstruction to the bile ducts was plain enough, the question was as to the cause. Without any definite data to sustain us, we came to the conclusion that his condition was due either to carcinoma of the biliary passages, or to their obstruction by gall-stones. In speaking of some of the causes of obstructive jaundice, you will remember that I referred also to hydatid cysts, and round worms as very rare causes of such a condition, so rare that one would hardly be justified in making such a diagnosis except by the most positive exclusion of the other causes of jaundice. You will also remember that I suggested that the question could be solved by means of the exploratory needle and by laparotomy, but that the patient was so ill that these means were scarcely justifiable. Moreover, his blood was so seriously poisoned by the retention of bile, that on this account an operation would not be justifiable, since the patient would probably die of hemorrhage, either from a copious escape of blood at once or from constant oozing. The fatal termination resulted four or five days afterwards, his stupor deepening into coma and death.

The autopsy showed a very remarkable state of affairs. There was no disease outside of the ducts or liver. The tissues and structures showed simply an impregnation by bile. The large common duct was very much enlarged, exceeding the size of my thumb. It could be readily distinguished from the vein alongside of it, and was easily isolated. There was not a sufficient amount

of adhesions to cause any doubt as to the relation of the parts. The gall-bladder was enlarged, and distended with a mucoid fluid. That portion of the duct extending from the cystic duct to the duodenum was patent. The biliary ducts in the liver were dilated, and just above the entrance of the hepatic duct into the liver there was a large dilatation, apparently of the duct, but which I believe to be a cyst which had ruptured into the duct and in that way occluded the canal with the hydatid membranes. The duct itself was not the seat of any disease. This cyst, hydatid in nature, was large enough to hold a good-sized lemon. There was a second cyst whose contents still remained in the periphery of the left lobe of the liver. It had nothing to do with the cause of the jaundice. The contents of the larger cyst, by causing an obstruction of the duct, had given rise to the jaundiced condition of the patient. These large shreds of membrane, which I here show you, could never have passed through the duct into the duodenum. Nature made an effort to accomplish this, but was not able to sufficiently distend the duct.

The question now is, could this man have been relieved by an operation? I think he could, because of the size of the duct. There were so few adhesions that access to the ducts could have readily been had and a cholecystotomy performed relieving the obstruction, while a rubber tube would have drained the cyst, and the patient would probably have recovered. Of course there would have remained the other cyst, which would gradually have increased in size; and, knowing this history of the patient, a second operation could have been performed, either by laparotomy or aspiration. The patient just missed his chance of recovery by not entering the hospital a few weeks earlier.

The lesson we learn from this case is, that an exploratory operation in obscure cases of obstructive jaundice is always advisable. Opening the gall-bladder simply, would not have been of service here. The mucoid contents of the bladder simply indicate that there is an obstruction of the cystic duct, and that the gall-bladder has merely become a large retention cyst, the removal of which would not cure the disease. The fact that this man died of an hydatid cyst of the liver, the contents of which obstructed the duct, would never justify you in making such a diagnosis. It is very probable that never again will you see an instance of such

¹Delivered at the Philadelphia Hospital.

a condition, unless you settle in some country where it is common, as in Australia or Iceland or perhaps northern Minnesota, in our own land.

Pleurisy with Tuberculosis.

I wish now to call your attention to this patient. He is a slight, delicate-looking young man, 20 years old, a stone-cutter by occupation, who has been much exposed to changes in the weather. His ailments in childhood were not unusual, and his habits were not bad. He has used alcohol moderately. There is nothing peculiar in his family history, and we can find but very little about his residence and his associates—an important point to learn in suspected cases of developing tubercular disease. On the thirteenth of July after doing some heavy lifting, he noticed that he was short of breath. About the same time his work was such that he was frequently chilled and wet through. This shortness in the breathing continued, and was attended with a slight amount of coughing, only noticed on stooping. At the time this cough was most severe, he had a pain in his right side, but none in the left. On July 29, he was obliged to quit work, and was admitted to the hospital on August 12, exhibiting these symptoms: great dyspnoea, especially on walking about; respiration increased somewhat, but not greatly—28 to 34 per minute—cough and fever with great depression and prostration.

Here we have a case of a sub-acute illness with the symptoms pointing to some disease of the chest, occurring in a lad who has been so much exposed that we are led to suspect some form of pulmonary disease. Throughout the whole time since his admission he has had fever, and he has also lost flesh to a moderate degree, while he suffers with night-sweats and great prostration. Note, therefore, in the first place, the gradual onset of the disease, and in the second place, the lung symptoms associated with fever, emaciation—not rapid—and night-sweats. We will now proceed to an examination of his chest.

His respirations, you will notice, are hurried, and there is some flatness on the right side of his chest, with fulness on the left. The lower part of the right side expands, while the left base scarcely moves. On palpation there is found a diminution in the vocal fremitus on the left side, with normal fremitus upon the right. On percussion,

there is found dulness in the left second interspace, increasing to flatness further down, with tympany over the gastric area. The right side shows normal pulmonary resonance. On auscultation, I find diminution in the breath sounds upon the left side, with a somewhat exaggerated respiratory murmur on the right side, puerile or bronchovesicular in character. We have here a unilateral increase in the size of the chest, which means a unilateral increase in the amount of its contents. Unilateral diminution in movement shows a diminution in the amount of air there, which is also shown by the absence of normal pulmonary resonance. We are able, therefore, from inspection, to eliminate the fact that this enlargement is due to some enlargement of the lung, since, if due to compensatory emphysema—the usual cause of such an enlargement—there would be an increased amount of air, with exaggerated movement of the chest. That it is in the pleura that we have this increase in the contents, is shown by palpation and in the absence of vocal fremitus. There is something here that prevents the vibration of the voice from reaching the hand. Without doubt, this is a pleural effusion.

We must determine now the character of this fluid, and the cause of the effusion. As to the nature of the fluid, we may learn this either by direct exploration, or by a further study of the physical signs. Serum, blood or pus may be present in the chest. If serum, the symptoms are such as you see present in this case; there are no physical signs by which we can determine the presence of blood from serum; but when pus is present we have the sign of Bacilli to rely on, that is, no transmission of the whispered voice. In this case a hypodermic needle has been used for diagnostic purposes, and a small amount of serum has been withdrawn. Such a procedure must be carried out with care, since, unless done antisceptically, empyema will result.

There are general and local causes of pleural effusion. The heart, kidneys and blood are free from disease, and thus the usual causes being eliminated, we are driven back to the pleura itself in investigating as to the cause of this condition. We learn that the disease developed insidiously, with shortness of breath, slight cough and fever.

Therefore, associated with this local symptom of serum in the chest, we have fever, emaciation and night-sweats.

As an aid to our discussion let us ex-

amine a similar case of effusion, with different general symptoms. Here is another well-appearing man who presents similar local symptoms and signs, which are not so marked as in this young man. There is lessened movement of his chest, slight fulness of the intercostal spaces, dulness on percussion, not very high laterally, and posteriorly beginning at the middle of the scapula and ending in flatness at the base, with absent breathing sounds. He has also slight shortness of breath, and slight pain; but otherwise he is not suffering at all. At times you will see such cases presenting no general symptoms; and if you are not careful to examine the patient all over from time to time, such a condition will often be overlooked. It is a very common thing to find cases of pleural effusion developed insidiously. As soon as the effusion takes place all pain disappears, and the patient may not lay much stress upon the slight pain present at first and lasting only a short period. Why, now, is there such a difference in the symptoms of these two cases? It must be due to a difference in the cause or to a difference in the character of the effusion. It is not due to the amount of fluid present; for this will have no effect upon the fever, emaciation and sweats.

Pleurisy with fever, when of long duration, is due to associated complications, to the peculiar nature of the effusion (purulent), or to a difference in the cause. Is there any specific process which is the cause of this condition in this lad? Is there any septic process here? A woman after confinement will have a chill, irregular fever, and a stitch in her side, a true pleurisy due to septicemia or pyemia. There is nothing like this here. There is no suppuration in any part of the body which would cause septicemia. In cases of tuberculosis, which is another true microbial process, pleurisy, like inflammation of other membranes, is apt to occur. Indeed many believe that pleurisy never occurs except of tubercular or septic origin. Is tuberculosis present in this instance? We have not been able in one examination to demonstrate the presence of bacilli in the sputum or serous exudation. It does not follow that tuberculosis is absent. The micro-organisms may be present in small numbers, and be detected only after frequent examinations. There is, however, undoubted tuberculosis at the apex of the right lung. The physical signs—upon which time will not permit me to detail—

show this. The co-existence of this process in the right lung, with the fever, emaciation and sweats, are sufficient to warrant a diagnosis of tuberculosis.

The points to remember here are: the oftentimes insidious onset of pleurisy and its association with tuberculosis.

The treatment of the case varies with the cause. In acute pleurisy with effusion the saline treatment is of service; but in this kind, with fever and prostration, we do not use such depletive measures. The fact that tuberculosis is associated with pleurisy does not compel us to make a fatal prognosis. Tubercular disease of serous surfaces is the most favorable form of the disease. Aspiration, or removal of the fluid in tubercular peritonitis, will often cure the disease; and so aspiration of the pleural sac will favor the cure of such cases. This should be done frequently.

COMMUNICATIONS.

CASE OF OVARIOTOMY.¹

BY JOHN L. ATLEE, M. D.,
CHATTANOOGA, TENN.

Fifty years ago, when the operation for the removal of an ovarian cyst was revived by the late Dr. John L. Atlee, of Pennsylvania, and his associates, and for years and years following their renewal of the operation, the man who was courageous and heroic enough to enter the abdominal cavity with the knife was indeed noted and eminent among his confrères. To-day, with the advance in science and surgery, the doctor who has *not* successfully performed a laparotomy is the one who stands noted among his colleagues. Indeed, so very common are the reports of the operations of laparotomy and ovariotomy, that I would not venture to report this case were it not that there are several features connected with it that I have never known to occur before, and which I think will make the report of interest.

In May, 1889, I was called to see Mrs. L., white, 21 years old, married two years, without issue. She menstruated at the age of fourteen, and continued regular. Directly after her marriage, she noticed an en-

¹ Read before the Tri-State Medical Association, October 16, 1890.

largement in her left side, which gradually increased until it became so considerable as to impede locomotion and interfere with respiration. Because of this preying upon the vital sources of the system, and because of the indigestion, which was a prominent symptom, I found my patient greatly emaciated and weak. Upon examination, I diagnosed a multilocular cyst of the left ovary, and appointed May 15, two weeks later, as the time for operation. On that day I found her with a pulse of 120, her temperature 103° , and her respiration very rapid, and with these symptoms I thought it best not to operate. Upon her earnest solicitation, and that of her family, for some immediate relief, I performed paracentesis, and drew off thirty pounds of a thick, coffee-colored fluid, albuminous in character, and presenting under the microscope the ovarian germ of Drysdale. The operation gave her great relief. On May 25, ten days afterwards, I operated, being assisted by my son, Dr. James H. Atlee, Dr. Force, of Calhoun, and Dr. Richardson, of Charleston, Tennessee, and removed by abdominal section a tumor weighing forty-five pounds, which in every way confirmed my diagnosis. The pedicle, which was short and thick, I secured with two ligatures, before severing the tumor: using one for each half of the pedicle. The ligatures used were Snowden's iron-dyed black silk ligatures, No. 14. After the tumor was severed, there was a profuse hemorrhage, due to the retraction of the artery from the ligature. Another ligature, cast around the stump, secured the artery and stopped the hemorrhage. After thoroughly cleansing the abdominal cavity with warm carbolized water, the divided edges of the peritoneum were brought together with cat-gut ligature, and the abdominal wound was closed with the iron-dyed ligature, with a drainage-tube of rubber in the lower portion.

The case progressed without a single unfavorable symptom, with adhesion by first intention. About the eighth week after the operation, symptoms of cystitis appeared, with difficult and painful urination. The urethra was found to be closed with an object, which, upon removal, was identified as one of the ligatures used in ligating the pedicle. Following the removal of the ligature from the urethra there was a flow of bloody pus and urine, with a subsidence of all disagreeable symptoms and a speedy recovery.

About the sixth month after the operation cystitis re-appeared, and gradually increased in severity until the second ligature was passed from the bladder. This ligature I have now in my possession. The passage of this ligature was, in a like manner, followed by a flow of bloody pus and urine, and the rapid abatement of all symptoms of cystitis. Since then my patient has remained in perfect health, and is now a stout, strong, healthy woman, busy with the duties of her household.

The interesting feature of this case is the passage of the ligatures from the peritoneal cavity, through the peritoneum and the walls of the bladder and through the urethra. I have no recollection of an ovariotomy that has progressed just as this one did, with the passage of the ligatures through the walls of the bladder and urethra, at the same time occasioning no inflammation of the peritoneum, no escape of urine into the peritoneal cavity, and no cystitis sufficient to endanger the life of the patient.

219 Oak Street.

CARCINOMA OF THE GALL-BLADDER AND LIVER FROM BILIARY CALCULI.

BY MARIA MITCHELL VINTON, A. M., M. D.,
INSTRUCTOR IN PHYSICAL DIAGNOSIS, WOMAN'S MEDICAL COLLEGE, NEW YORK.

Carcinoma of the liver is, in three-fourths of the cases, secondary, the primary growth occurring in the stomach, intestines, pancreas or mesentery, and being communicated to the liver either by direct extension or by metastasis. Carcinoma of the gall-bladder, extending into the liver, is comparatively rare. Such a carcinoma is the result of the formation of a mass of gall-stones in the gall-bladder, this causing attacks of inflammation and adhesions forming between the liver and the gall-bladder, which become the seat of carcinomatous growth. Carcinoma of the liver is usually nodular, forming a more or less spherical mass protruding from the surface of the liver and changing its shape. When extending from the gall-bladder, it infiltrates the liver substance, giving an even enlargement. The usual form of growth is of the encephaloid type, cell-nests predominating over fibrous tissue.

The following report describes a case of

carcinoma of the gall-bladder extending into the liver, occurring with no previous attacks of biliary colic or other symptoms pointing to the formation of a mass of gallstones. That a mass of over twelve hundred stones, some of them of considerable size, could form without giving rise to symptoms sufficient to attract the patient's attention, seems almost impossible.

E. M. was seventy-nine years and eleven months old at the time of her death. The family history was very incomplete. Two sisters had died of cancer—in what situation was not known. The patient had had several children who, at the time of her death, were all living and healthy. She could give no history of any past illness more severe than constipation, and had never had biliary colic. She was seen first at the Dispensary of the New York Infirmary for Women and Children, July 2, 1889. She then complained of having had, for about two weeks, nausea, pain and fulness in the stomach after eating. Her tongue was coated and her bowels constipated. Simple gastric remedies relieved her, and no physical examination was made. She came again January 2, 1890. She then complained of the same symptoms as before, but had also a lancinating pain over the liver running through to the back. She stated that she had had this pain for three months, and had not been feeling as well as usual for a year. Examination of the abdomen showed the liver to be enlarged, projecting a finger's breadth and a half below the border of the ribs in the nipple line. The edge was smooth and hard. Below the liver, and attached to it, but separated from it by a sulcus, was a mass the size of a man's fist, which felt distinctly nodular through the lax and thin abdominal wall, and of stony hardness. This mass was tender on pressure, and was the seat of the lancinating pain complained of. The other abdominal organs were normal, as were also the heart and lungs. The urine was stated to be scanty and contained no albumin, but a few granular casts. The patient was thin, but not cachectic. She had simply the look of poor nutrition, so familiar at dispensaries.

A diagnosis was made of carcinoma of the liver, probably secondary to a deposit in the pyloric end of the stomach or duodenum. The patient continued to visit the clinic up to about February 1, when she became unable to leave her house. She frequently located her pain in the epigastrium, instead of over the liver. She had no vomiting and

no jaundice. The bowels were constipated throughout her illness. The stools contained, at times, blood and mucus, and were never clay colored.

By February 8, the patient had acquired the cancerous cachexia, was weak and much emaciated. The pain had increased much. On February 13, vomiting occurred for the first time.

The vomited matter contained no blood. The urine showed a small amount of albumin, but no bile pigment. It deposited a very heavy precipitate of urates. The tumor had increased much in size, and was extremely tender. It seemed now to consist of two distinct masses, separated by a deep sulcus. The spleen was of normal size.

On February 20, the patient's temperature was 100.4°. The patient's mind had not been clear for several days, she was delirious during the previous night, and remained in a semi-conscious state. She complained of pain in the epigastrium, but not over the liver. The tumor had been enlarging rapidly, and now reached the umbilicus, and the liver extended down over the lower mass.

February 21, she had marked fever; the skin was slightly jaundiced; but there was now no complaint of pain. Her urine had a specific gravity of 1.029, was dark brown in color, giving a marked yellow froth on shaking, and a play of colors with the sulphuric acid and sugar test. It contained a trace of albumin.

February 22, the temperature was 99.8°, and the woman had severe pain over the liver and was deeply jaundiced.

On February 23, she had a succession of chills, lasting all day, and towards evening became very feverish. She suffered intense pain, so that she cried out and flexed the legs on the abdomen. On February 24, the pain had passed off, but she remained semi-conscious, and died early on the morning of February 25.

An autopsy was made by Dr. Walter Vaught. The lungs showed a few milliary tubercles at the left apex and firm pleuritic adhesions on the right. The heart was normal. The pancreas, spleen and kidneys were normal. The stomach—the suspected seat of primary carcinoma—was normal. The jejunum and ileum were normal. The colon was crowded with hard scybala masses of feces, but entirely free from carcinoma. The right lobe of the liver projected two finger breadths below the border of the ribs. The

anterior surface of the right lobe over the region of the gall-bladder was of stony hardness; the left lobe was normal in size. Projecting from the lower surface, in the location of an enlarged gall-bladder, was a hard nodular mass, the size of a man's fist, firmly adherent over its entire upper surface to the liver. In cutting into this mass, the gall-bladder was opened and showed the mass to be composed of gall-stones of all sizes, firmly pressed together and forming a single mass, which was not easily broken up until the incision in the gall-bladder had been lengthened. After removing the stones, the walls of the gall-bladder were seen to be infiltrated with cancerous material, which had extended into the substance of the liver. The mass in the liver formed a patch about the size of the palm of a man's hand, varying from one-quarter inch to one and one-half inches in thickness, smooth and not nodular. The substance was firm and of a white or pinkish color on section, contrasting with the deep color of the congested liver tissue. All the bile ducts were in a state of suppuration, and filled with a mixture of pus and bile. The mucous membrane of the gall-bladder was ulcerated in many places. One ulcer had perforated the wall of the duodenum and formed an opening the size of a five-cent piece. In the cystic duct, at its junction with the hepatic, was found a gall-stone, the size of a filbert and firmly impacted. This entirely prevented the passage of bile into the intestine, and as jaundice did not occur until four days before death, the stone must at that time have reached its position at the junction of the ducts, and the severe colic on the day before death was probably due to its attempt to pass down the duct. That the duct must have been partly occluded four days before death is shown by the beginning of jaundice at that time. This caused the suppuration of the bile ducts and the fever and delirium of the last few days. Several inches of the duodenum were included in the cancerous mass, all being firmly matted together. The lumen of the intestine was in one place contracted so as to admit only the tip of the finger. A section of the cancerous mass in the liver showed cell-nests filled with large cells and separated by a more or less scanty stroma. The deposit had occurred by infiltration directly from the adherent gall-bladder.

The gall-stones were of three distinct kinds. There were about a dozen the size of a filbert and of a whitish color. The re-

mainder were of a dark-gray with a yellowish surface easily flaked off by the nail. There were four hundred, varying in size from that of a small French pea to that of a hazel nut. Besides these, there were a large number of smaller ones, like fine granular sand. Of these eight hundred were counted and many were lost. They appeared to consist of cholesterol, variously colored by bile pigment and were very hard.

While years must have been occupied in the formation of this mass of gall-stones, they had produced no biliary colic, and had never caused sufficient irritation to attract the attention of the patient or of any physician, until the growth of the cancer, five months before death.

160 East Thirty-sixth street, New York.

PERISCOPE.

Diabetic Coma.

The *Lancet*, November 8, 1890, says: Dr. R. Schmitz, of Neuenahr, in Germany, has had a large experience in the treatment of diabetes, and therefore his opinion on diabetic coma is to be regarded with interest. He states that under the term "diabetic coma" two distinct conditions are to be recognized, both of them serious and somewhat similar at first glance, but entirely different as regards their pathology, and requiring different modes of treatment. The first is simple collapse, with coma; the second is an auto-infection, and is diabetic coma properly so-called. By most observers it is ascribed to the condition of acetonuria. The collapse is the result of weakened heart power, brought about by an excess of sugar in the blood causing a general degeneration of the muscular tissue throughout the body, in which the heart participates. During collapse there is marked cyanosis, the respiration is quickened, the pulse is generally about 68 to 72 a minute, the temperature is not raised, the heart's beat is almost imperceptible and the first sound can scarcely be heard. The treatment of this condition, according to circumstances, is prophylactic or active. If the cardiac degeneration is recognized before collapse occurs, the patient must be particularly warned against any violent exertion, cardiac depressants (such as the bromides or antipyrin) must be avoided, but in addition to the ordinary diabetic treatment

a little alcohol should be added to the diet, and the patient should be out in the fresh air as much as possible. During the actual attack stimulants must be employed, of which the best are black coffee, castor oil and hypodermic injections of camphor and musk. In true diabetic coma there are generally well-marked prodromal symptoms—complete loss of appetite, somnolence with unrefreshing sleep, etc. The tongue becomes coated and dry, the breath very fetid. When coma sets in there is fever (100° to 102° F.), respiration and pulse are quickened, the former rising to 45 and the latter to 130 a minute. There is much epigastric pain, but there is no tenderness. The diagnosis is easy and is based on the epigastric pain and rise of temperature (both of which are absent in the first form), on a negative result from an examination of the heart, and on the rapidity of the onset of the coma. The treatment is directed to the removal of the toxic material, which Dr. Schmitz considers accumulates in the intestine. He consequently advises frequent doses of castor oil, even if diarrhoea be already present. He orders half an ounce every hour until a full evacuation is produced, and states that he has had excellent results from this method of treatment.

Salinaphthol in Rheumatism.

Salinaphthol is a combination of salicylic acid with naphthol, and is analogous to salol, which is a combination of the same acid with phenol. It is insoluble in water, and has neither odor nor taste.

In a review of salinaphthol, the *Journal de Médecine*, November 9, 1890, says that the remedy seems to act as well as salol in acute articular rheumatism, and it is well borne. In the dose of five and one-half to seven and one-half grains, repeated four times a day, it produces neither headache, ringing in the ears, nor any symptom of poisoning, even after prolonged use. Kober claims that it is separated into salicylic acid and naphthol by the pancreatic juice and by the ferment which the intestinal mucous membrane secretes, and that this separation is evidenced by the presence of salicylic acid in the urine. Lépine denies that the separation takes place from the action of the intestinal juice, and admits such separation only as the result of the action of the pancreatic juice. There is here room for fresh experiments to examine whether or

not the pancreatic juice really has this action, and if it is not the same for salinaphthol as for salol, which is decomposed not only by the pancreatic juice, but also by the alkaline constituents of the blood.

Salinaphthol should be less harmful than salol, since naphthol is less toxic than phenol. But, up to the present time, clinical experience does not seem to have established the efficaciousness of this medicament.

Railway Readers.

The *Lancet*, November 8, 1890, says: a contemporary recently stated that a French medical practitioner has been collecting statistics with regard to those of his patients who complain of nervous affections, and has come to the conclusion that all the evil proceeds from the practice of reading in the train. We can hardly accept this view. Many thousands of business men have no other time than the half or three-quarters of an hour to learn the state of the markets, the last moves in the politics of the great Powers, general news, and the various items that make up the contents of a morning or evening paper, and it would hardly appear that the duration of the cause is sufficient to produce the effects assigned to it, whilst the majority of people travel too rarely and read to little to suffer at all. Our contemporary suggests that a paper with specially large type should be printed for the use of travelers by rail. We fear it would prove of little service. We suggest that all carriages should, as a general rule, be supplied with a better light.

Ergotin in Gleet.

In the Polish periodical, *Wiadomosci Lekarskie*, No. 8, 1890, p. 250, Dr. A. Roicki highly eulogizes ergotin as a means for rapidly curing chronic gonorrhœa. The drug should be simultaneously used, both internally, in pills, and locally, after the following formula.

R Ergotini 0.3 grammes
Aqua destillata 300.0 grammes
M. D. S. Inject a syringeful into the urethra several times a day.

The injections are said to be tolerated perfectly well. The same treatment proves useful in cases of urethral hemorrhage.

xiii
ac-
aph-
not
the
than
sph-
ical
shed

Dec. 27, 1890.

Editorial.

733

THE
MEDICAL AND SURGICAL
REPORTER.
—
ISSUED EVERY SATURDAY.

CHARLES W. DULLES, M.D.,
EDITOR AND PUBLISHER.

N. E. Cor. 13th and Walnut Streets,
P. O. Box 843. Philadelphia, Pa.

SUGGESTIONS TO SUBSCRIBERS:

See that your address-label gives the date to which your subscription is paid.

In requesting a change of address, give the old address as well as the new one.

If your REPORTER does not reach you promptly and regularly, notify the publisher *at once*, so that the cause may be discovered and corrected.

SUGGESTIONS TO CONTRIBUTORS AND CORRESPONDENTS:

Write in ink.

Write on one side of paper only.

Write on paper of the size usually used for letters.

Make a few paragraphs as possible. Punctuate carefully. Do not abbreviate or omit words like "the" and "a," or "an."

Make communications as short as possible.

NEVER MAIL A MANUSCRIPT! Try to get an envelope or wrapper which will fit it.

When it is desired to call our attention to something in a newspaper, mark the passage boldly with a colored pencil, and write on the wrapper "Marked copy." Unless this is done, newspapers are not looked at.

The Editor will be glad to get medical news, but it is important that brevity and actual interest shall characterize communications intended for publication.

THE END OF THE YEAR.

A PRIVATE WORD WITH OUR READERS.

During the past year the columns of the REPORTER have contained—as will be seen by looking over the Index and the list of Contributors—a large number of scientific papers of great value from men deservedly eminent in the profession and from many others of less extensive reputation, but who are careful observers and skillful practitioners. The practical value of the papers submitted for publication has always been regarded as the best criterion of their usefulness. Especial attention has been bestowed upon the feature of Clinical Lectures; because it is believed that many readers like to hear what they would be hearing if they were again in the schools where the art of medicine is taught. The

Department of Correspondence has covered a large number of American and foreign cities, and has included short communications from readers of the REPORTER in every part of the world. The Selections, Translations and Abstracts have been made with great care, and their source has been fully stated, so that proper credit should be given to those of our contemporaries from which they have been taken, and so that any reader might go at once to the original if he wished. The Editorials and Book Reviews, written by the Editor and a number of men especially competent in various departments of medicine, have always been prepared with an eye single to the good of the reader. No interest, no friendship has been allowed to stand in the way of honest criticism; and no praise has been insincerely uttered.

The purpose of the REPORTER is to be a journal wholly and fearlessly devoted to the advancement of the medical profession, and an alert and conscientious friend to every one who consults its pages. This purpose will shape its policy in the year upon which we are about to enter, and in which we hope all the readers of its pages may be abundantly prospered.

HYSTERIA IN MALES.

In a communication to the *Archives de Médecine*, August, 1890, Dr. A. Souques reports thirteen cases of hysteria in men which he has seen in the Broussais Hospital. From an abstract of this paper in the *Bulletin Médical*, October 1, 1890, we learn that Souques thinks that hysteria in the male is by no means uncommon, that it is encountered frequently in ordinary hospital practice, and that it may be more frequent in the lower classes than hysteria in women. The patients he observed were from twenty to fifty years of age, and were nearly all working men, while some of them were in actual destitution. According to Charcot, such persons are more subject than

others to the causes which sometimes give rise to hysteria, such as traumatism, intoxications and infectious diseases. Souques found the cause of hysteria in his cases to be lead poisoning eleven times, traumatism three times and once each syphilis, dysentery and alcoholism. But there must also be an hereditary nervous tendency.

The symptomatology of hysteria in males is variable. Often the onset is marked by an apoplectiform attack, more often, perhaps, by vertigo, dazzling sensations and a disposition to faint. Vertigo appears to characterize hysteria in the male very much as convulsions do hysteria in women. The most frequent motor troubles of hysteria in males are paryses, while anaesthesia, limited by the median line of the body, are common. In general demeanor a female hysterical patient is usually gay and lively, while a male is likely to be sombre and gloomy. Hysteria in males is a relatively mild disorder; and usually it terminates favorably. The motor troubles are the most rebellious, and the prognosis in regard to them should be reserved. The patient will probably recover, but it cannot be said at what time. The possibility of relapses should be borne in mind.

The treatment of hysteria in males should be directed to both the hysteria and the provoking cause.

Those who feel interested in the subject of hysteria in men will be interested, we think, to know that Dr. Julius Dreschfeld has an excellent article, "On some of the Rarer Forms of Hysteria in Man" in the *Medical Chronicle*, October, 1890. Dr. Dreschfeld takes almost precisely the same view of the etiology of this disorder as does Souques—who appears to have adopted it from Charcot—and reports four cases which differ from those usually observed after shock and injury. The first two cases belong to the group known as spurious hydrophobia or pseudo-hydrophobia. They are similar to cases familiar to all who have made a thorough investigation of the litera-

ture of hydrophobia, and—aside from their intrinsic interest—would prove instructive to persons unfamiliar with the disorders which simulate hydrophobia.

The whole subject of male hysteria is one of great interest, and we think it is one in which there is still room for much investigation. The name is unfortunate; but until a better one is suggested it will have to serve to indicate what is now generally—though vaguely—understood by it.

A KOCH INSTITUTE IN NEW YORK.

In striking contrast to the attitude of caution which all scientific medical men are observing towards the Koch remedy for tuberculosis, it was announced on December 21, that a "New York Bacteriological Institute" had been incorporated for "the study and gratuitous treatment of contagious diseases, comprising a Pasteur and Koch department for the treatment of hydrophobia and tuberculosis." In the list of Trustees, the observant reader will not be surprised to find the names of Drs. Gibier and Lautard.

BOOK REVIEWS.

[Any book reviewed in these columns may be obtained upon receipt of price, from the office of the *REPORTER*.]

CYCLOPÆDIA OF THE DISEASES OF CHILDREN. EDITED BY JOHN M. KEATING, M. D., Volume iv. Large 8vo, pp. xii, 1128. Philadelphia: J. B. Lippincott Company, 1890. Sold by subscription.

It is sometimes hard to find a new thing to say about an admirable book, especially when the space which can be given to a single review is so limited as it is in the *REPORTER*. It is no small praise to say that the fourth volume of Keating's *Cyclopædia of the Diseases of Children* maintains the high standard of those which preceded it. This volume concludes the work; and the whole Cyclopædia stands a monument of medical, editorial and publishing enterprise and skill, of which American medical men may justly be proud. The present volume contains articles on the ear, the eye, hygiene and diseases of the nervous system. Among them, it would be difficult to select those which are most deserving of commendation; but, in noticing this last volume, we think it is not invidious to call especial attention to the admirable article on physical development by the editor, Dr. Keating, and Dr. J. K. Young.

We call attention to this particular article, because it is a pleasure to add this tribute to the ability which Dr. Keating has displayed in preparing this unusually valuable work.

KO

A g
propos
tuber
shows
cal Co
he had
power
bacilli
the bo
guinea
ceptib
influen
the in
that i
tuber
process
still."

This
citemen
the m
Koch
"sub
gress
which
same."

Aft
to be
the "
and c
upon
newsp
patien
with i
them."

Pro
ports,
Medi
13, 1
was i
paper

In
from
reme
admin
dersta
to pr
be ca
diagr
of it
tuber

SPECIAL ARTICLE.

KOCH'S REMEDY FOR TUBERCULOSIS.

A RESUMÉ OF ITS HISTORY.

A glance over the history of the remedy proposed by Koch for the treatment of tuberculosis (in the strict sense of this term) shows that at the Tenth International Medical Congress, last August, he announced that he had discovered a substance which has the power of preventing the growth of tubercle bacilli, not only in the test-tube, but also in the body of an animal. He also said that guinea-pigs, which are extraordinarily susceptible to tuberculosis, "if exposed to the influence of this substance cease to react to the inoculation of tuberculosis virus, and that in guinea-pigs suffering from *general tuberculosis even to a high degree, the morbid process can be brought completely to a standstill.*"

This announcement created intense excitement and great enthusiasm, and aroused the most sanguine hopes that before long Koch would disclose the character of the "substance" which could arrest the progress of tuberculosis in guinea-pigs, and which he evidently expected would do the same in human beings.

After the Congress had adjourned it came to be believed that Koch had given some of the "substance" to certain practitioners and clinicians of Berlin for experimentation upon their patients, and before long the newspapers began to contain statements that patients with consumption were being treated with Koch's remedy, and then that some of them were cured.

Prompted probably by these imperfect reports, Koch, in a special issue of the *Deutsche Medicinische Wochenschrift*, dated November 13, 1890, made an announcement which was immediately sent by telegraph to newspapers in various parts of the world.

In this communication Koch refrained from disclosing the nature of his proposed remedy, but indicated what he thought its administration would accomplish. To understand what this communication seemed to promise, certain of its expressions must be carefully weighed. After describing the diagnostic value he attributed to it because of its effect upon persons believed to be tuberculous, Koch said: "Very much more

important than the significance which the remedy (*Mittel*) has for diagnostic purposes, is its curative action (*Heilwirkung*)."¹ He then described its employment to cure two cases of lupus, and in the treatment of tuberculosis of the glands, bones and joints, and says "the success was the same as in lupus, rapid recovery (*Heilung*) in fresh and light cases, slowly progressing improvement in severe cases." When speaking of the effect of his substance on consumptives, he said: "The action of the remedy expressed itself in consumptives in general in such a manner that the cough and expectoration usually increased somewhat after the first injections, then, however, became less and less, so that in the more favorable cases they finally disappeared entirely." With loosening of the sputum and the disappearance of bacilli in it, until it too ceased, he says that night-sweats ceased, the appearance improved, and the patients gained in weight. As to the ultimate results, he said: "Patients treated in the incipient stage of phthisis were all freed from every symptom of disease in the course of from four to six weeks, so that they could be regarded as cured." He said also: "Patients with not too large cavities were markedly improved and almost cured. According to those experiences I would assume that *incipient phthisis may be cured with certainty* (*Sicherheit*) by the remedy. This may be partially true also of cases which have not progressed too far."

Koch recognized the limitation necessary to his statements on account of the short period during which the cases he thought cured had been under observation, but he said it might well be assumed that possible relapses would be "as easily and quickly cured as the first attack."

The publication of this announcement produced the most extravagant outburst of applause and raised the most boundless expectations that a cure for consumption had been discovered. The daily papers gave the greatest prominence to the doings and sayings of all connected with the experimental application of the substance to which the newspapers gave the name of "lymph,"¹ and patients and medical men flocked to Berlin to get or to have something from Koch and those to whom he entrusted it.

¹ The word "lymph" is no doubt a misnomer, but is used here as a convenient and now recognized term for what Koch called a "remedy" (*Heilmittel*).

The first full and authentic accounts of what was being done with the "substance" came in the *Deutsche Medicinische Wochenschrift*, of November 20, 1890, which contains reports of the experiences of Fräntzel and Runkwitz, Levy, Köhler and Westphal.

Fräntzel began to use the "lymph" on September 13, and he and Runkwitz give accounts of four cases of far advanced phthisis, none of which were benefited and two of which ended in death before the report was made; of eight cases of moderately advanced phthisis, in which some improvement was noted.

Levy gives an account of the first three cases of surgical tuberculosis treated by Koch's method—the first a case of lupus injected on October 10.

Köhler and Westphal began their treatment on October 11, on eight patients with tubercular disease of the bones and skin, and four patients without tuberculosis. Their account is most full and complete, but they conclude by saying it is only a contribution of facts, and they do not venture to pronounce a judgment in regard to the value of the remedy.

Von Bergmann's communication was the address he delivered before the Free Surgical Association of Berlin, November 16, 1890. In it he gave the results of his experience and mentions the following application of Koch's remedy, exhibiting the patients to the assembly.

LUPUS—13 CASES.

No. of Patients	No. of Inoc's.	Dates.
1	5	Nov. 6, 8, 10, 12, 14
1		" 6, 7, etc.
1	1	" 6
5	1	" 16, A. M.
5	1	" 16, Ev'g

CERVICAL ADENITIS—3 CASES.

1	5	Nov. 6, 8, 10, 12, 14
1	1	" 16, A. M.
1	1	" 16, Ev'g

JOINT TUBERCULOSIS—16 CASES.

3	5	Nov. 6, 8, 10, 12, 14
1	4	" 6, 10, 12, 14
1	1	" 6
4	1	" 16, A. M.
7	1	" 16, Ev'g

TUBERCULAR LARYNGITIS—4 CASES.

1	4	Nov. 10, 12, 14, 16
3	1	" 16, A. M.

He also showed a man who had been supposed to have tubercular laryngitis, but whom he now believed to have carcinoma of the larynx because he did not react, as was expected, to a single injection of Koch's lymph administered that morning.

It will be noted that only nine of these thirty-seven patients had been under treatment as long as ten days; one had been under treatment for only six days; fourteen had had their first injection on the day of the meeting, and thirteen were injected while the meeting was being held. Careful reading of the address shows that Bergmann felt encouraged to hope that the lymph would prove valuable both in the diagnosis and in the treatment of surgical forms of tuberculosis.

The first person to use Koch's fluid in France was Prof. Péan, who, on November 29, described his experiments in a lecture at the Hospital St. Louis, in Paris. He gave a brief account of his observations in fifteen cases and concluded with the statement that "the results do not permit any conclusion in regard to the curative value of the medicament." (*Bulletin Médical*, November 30, 1890.)

Cornil in a lecture at the Laennec Hospital, November 30, spoke of having administered the "lymph" to a few patients the day before (November 29) and ventured no opinion whatever in regard to its curative power. (*Bulletin Médical*, December 1, 1890.) In a second lecture (*Bulletin Médical*, December 10, 1890) he said that the reaction indicated by Koch had appeared in every case in which he had used the lymph, and this was especially marked in cases in which the tuberculous process was of relatively recent beginning and active. He found the lymph to produce albuminuria and hematuria.

At the Société Medicale des Hôpitaux, December 5, communications were made by MM. Ferrand, Cuffer and Thibierge, who had just returned from Berlin where they had studied Koch's method under the auspices of the Berlin medical men. Dr. Thibierge had given especial attention to its effect on lupus, and had carefully examined a number of patients who were said in Berlin to be cured. He said to the Society that in view of his own investigations he felt justified in stating that there had not been a single case of even apparent cure of lupus with Koch's "lymph." (*Bulletin Médical*, December 7, 1890.)

Dec.

M.
"lym
called
depress
able as
cases
the pro

M.
Rémor
cases
in whi
(Bulle

Dr.
Moder
seems
so far,
to obt
served
jection
the m
fied."
gestio
used
tuber
the sa
and a
eating

Dr.
Heba
ber 1
just c
injec
10, c
Berli

O
dress
(Wi
goin
He s
the l

T
Mea
claim
had
was
mus
not
ing
to h
effe
call
of g
cal
com

ce

M. Cuffer described the effect of the "lymph" on cases of lung disease, and called attention to the fact that the remedy depresses the heart and that it is not reliable as an aid to diagnosis because certain cases of tuberculosis have failed to exhibit the promised reaction after its injection.

M. Debove said that his prosector, M. Rémond, stated that he had seen fourteen cases of undoubted tuberculosis in Berlin, in which the reaction had failed to appear. (*Bulletin Médical*, December 7.)

Dr. Charles Talamon, in the *Médecine Moderne*, December 11, 1890, says: "It seems on reading all that has been published so far, as if the only object of pursuit were to obtain the general and local reaction observed by Koch as a consequence of his injection. The more intense this reaction, the more the experimenters seem to be satisfied." He also makes the interesting suggestion that Koch's "lymph" should be used to determine the real nature of bovine tuberculosis—which has been denied to be of the same nature as tuberculosis in mankind—and also to decide what animals are fit for eating and what are dangerous.

Dr. L. Lereboullet, editor of the *Gazette Hebdomadaire*, says in the issue of December 13, that a patient with hip-disease had just died in coma after receiving his second injection by Péan, and that by December 10, eleven cases of death had occurred in Berlin attributable to this treatment.

On November 24, Köhler made an address in the Hygienic Institute of Berlin (*Wiener Med. Presse*, November 30, 1890), going over his experiences for six weeks. He said absolutely nothing new or justifying the belief that the lymph cures anything.

The special correspondent of the *Wiener Med. Presse*, November 30, quotes the disclaimer of Fräntzel and Bergmann that they had obtained actual cures. Fräntzel said he was "only a tyro," Bergmann said: "We must stick to the point that as yet we have not seen a single case of thorough healing." On November 29, Gerhardt said to him, "Here we can only speak of early effects, but not of curative effects." He also calls attention to the unfortunate appearance of greed of gain in cases of many of the medical men who flock to Berlin, and distinctly contrasts what he calls "*Auri sacra fames*" with the old statement: "*Dat Galenus opes.*"

The *Berliner Klinische Wochenschrift*, December 1, 1890, contains a communication by Israel in regard to the anatomical con-

ditions of two cases of joint tuberculosis treated with Koch's remedy. One specimen was taken from the ankle-joint of a patient who had received thirteen injections in the last eight days before the examination was made. The other patient had received seven injections, the last on November 19. One of these cases gave no anatomical evidence of being tuberculous, and the other was very far from conclusive.

The *Münchener Medizinische Wochenschrift*, December 2, contains an interesting article by Grünwald, illustrated, showing the effect of the method upon some of the cases of laryngeal tuberculosis treated by Bergmann already referred to. It also contains a summary of the recent publications in regard to Koch's method. Lublinski thinks that the remedy acts favorably upon laryngeal tuberculosis, although he admits that not a single case of real healing has been observed so far, and lays greatest stress upon the value of the lymph as a means of diagnosis.

The *Wiener Medizinische Wochenschrift*, November 30, contains a description of a chemical and microscopical analysis of the fluid of Koch by the Drs. Jolles, by which it appeared that the active principles were certain albuminoids, probably of the group of tox albumin or enzymes.

The *Berliner Klinische Wochenschrift*, December 8, contains an account of the proceedings before the Prussian Legislature in regard to the attitude of the Government towards the preparation and distribution of Koch's material, when Dr. Von Gossler, the minister of public worship (Cultus Minister) announced that the Government would assume control of the matter, and gave, as the chief reason for maintaining secrecy about its composition, the fact that at an interview with him on November 7, "in the presence of two Councillors of State, it was found that Koch was not able (*im Stande*) to give so exact a description of his process as to insure the preparation of an efficient remedy." Something he could tell but, in the opinion of Gossler, not enough to enable even the most skilful bacteriologist to follow his steps exactly and obtain the material he had obtained.

In the *Berliner Klinische Wochenschrift*, December 2, 1890, is a report of a lecture by Krause, on November 27, in regard to his experiences during two weeks and a half with Koch's method. This communication is also of an uncertain sort, expressing more

of hope than of confidence, and confessing that Krause has seen nothing which he could consider a cure.

The *Deutsche Medicinische Wochenschrift*, December 4, contains a communication by Rosenbach, of Breslau, in regard to his observations on fifty-six patients. Again we find the same unsatisfactory conclusions founded upon the short period of observation. Rosenbach distinctly states that he is very far from venturing any conclusions in regard to the therapeutic value of this material. A somewhat similar communication is published from Gerhardt's clinic by Carl von Noorden, in which—as is generally to be noticed in all the German communications—the cases are reported with thorough fidelity, and in sufficient detail to show the careful manner in which the observations have been made, and the candid scientific spirit of those who have made them; but it leads to the same entirely unsatisfactory result, and does not present any good ground for more than hope in regard to its efficiency.

The *Wiener Medizinische Presse*, December 7, 1890, contains an article by Jacksch, of Prague, who believes the remedy to be of a diagnostic value. He thinks that it is a very valuable expectorant in cases of pulmonary tuberculosis. He does not venture to express an opinion in regard to its value in the treatment of lupus.

[TO BE CONTINUED.]

NOTES AND COMMENTS.

Exact Dosage in the Cataphoretic Use of Drugs.

Dr. Frederick Peterson, in the *New York Medical Journal*, November 15, 1890, says that in a previous paper there are figured two cataphoretic electrodes devised for the anodal diffusion of drugs through the skin. The great drawback until the present moment, however, had been the difficulty of accurately regulating the amount of drug introduced. For this purpose rather complicated electrodes had hitherto been required, and even these had been unsatisfactory. He had recently found, however, that all difficulties are easily obviated by the use of a new and exceedingly simple method. Messrs. Waite & Bartlett have made for him a cataphoretic electrode of metal. Instead of covering it, as before, with sponge, the ordinary metal surface is

overlaid with a thin disc of platinum, and around the edge of this is placed a narrow rim of soft rubber. The drug to be used is put drop by drop upon a disc of ordinary tissue paper cut to fit the disc of platinum. Filtering paper or linen cloth may be used instead of tissue paper. A disc two or three centimetres in diameter will hold from one to four drops of the solution. When the medicated disc is placed upon the metal surface of the electrode, and the latter then applied to the skin, it is evident that there is a thin capillary layer of the drug in solution exposed to the cataphoretic power of the anode, between the electrode and the skin, and that the quantity of the drug used may be accurately estimated. The current is allowed to flow if desired until the medicated disc becomes perfectly dry. In this way one may introduce, he says, one or more drops of chloroform, methyl chloride, ether, ten-to-twenty-per-cent. solutions of cocaine, a one-per-cent. solution of helleborin, solutions of iodide of potassium, corrosive sublimate, aconitine—in fact, any drug to be employed in this manner.

To further simplify the method, Dr. Peterson has had medicated cataphoretic discs prepared by a pharmacist for use at any time, for the paper discs may be charged with any amount of a watery solution, and, the water being allowed to evaporate, they may be kept on hand indefinitely. It is only necessary to add two or three drops of water to the disc in administering the drug by electricity.

An apothecary has made for him and is prepared to supply any one with the following cataphoretic discs: Discs of menthol, 2 grains; of helleborin $\frac{1}{25}$ grain; of strychnine nitrate, $\frac{1}{2}$ grain; of iodol, 2 grains; of corrosive sublimate, $\frac{1}{6}$ grain; of cocaine hydrochloride, $\frac{1}{2}$ grain; of aconitine, $\frac{1}{4}$ grain; of potassium iodide, 4 grains; of mercury succinimide, $\frac{1}{4}$ grain; of lithium chloride, 4 grains.

Pleasant Laxative for Infants.

According to the *Revue Générale de Clin. et de Therapeutique*, March 5, 1890, Ferraud uses as a laxative for infants one or two dessert-spoonfuls of the following mixture in a cup of hot milk or weak tea:

R	Manna	3 vi
	Magnesia	
	Sulphur. loti	aa 3 iss
	Mellis	fz vi

Evidences of Criminal Poisoning.

In a paper read before the Medical Jurisprudence Society, of Philadelphia, November 11, 1890, Professor John J. Reese, M. D., read an interesting paper on the subject of "What Constitutes Reliable Evidence in Criminal Poisoning." The evidences he grouped under the heads of those derived from the symptoms before death; those derived from *post-mortem* appearances, especially as regarded the stomach; those obtained from chemical analysis; the evidences obtained by experimenting on living animals; and those derived from the circumstances of the case.

In reference to the evidences derived from the symptoms before death, he said it would be impossible to diagnosticate a case on such symptoms exclusively, as certain diseases presented similar symptoms. An exception might be made in poisoning with strong mineral acids, or with hydrocyanic acid. On the other hand, he had known cases of arsenical poisoning to be mistaken for cholera morbus. The second class of evidences, while strongly suggestive, like the first, could not, he said, be regarded as absolute proof, on account of the resemblance to pathological conditions found in certain diseases. It would not be safe, Dr. Reese said, to found a positive diagnosis on the pathological appearance of the lining of the stomach, as many cases of arsenical poisoning had occurred without leaving any evidence of its action on the lining of that organ.

In reference to evidences derived from chemical analysis, he said there were a number of poisons which could not be recognized by any known chemical tests, and other measures had, therefore, to be resorted to. The presence of a poison in the stomach or other organs did not necessarily mean that death was caused by the poison, as the latter might have been injected after death.

Where there was a failure to detect poisons by chemical analysis, it was necessary sometimes to resort to experiments upon living animals, such as dogs, cats and guinea-pigs. But often the most important evidences were those derived from the circumstances of the case, and, in the majority of cases, the evidence was mainly, if not exclusively, circumstantial. Rarely does it happen that there is direct proof of the administration of the poison.

In conclusion, Dr. Reese spoke of the various circumstances to be considered in a given case; such as the proven purchase and possession of the poison by an accused person; the exercise of an exclusive care of the victim during his illness by the accused; and the existence of a sufficient motive on the part of the accused.

Treatment of Delirium Tremens.

At the Congress of German Naturalists, September 15-19, 1890, Aufrecht, of Magdeburg, made some remarks upon the treatment of delirium tremens, which are reported in *Le Mercredi Medical*, November 5, 1890. Two questions, he said, present themselves: Can the disease be radically cured? Is it useful to continue to give alcohol to patients with delirium tremens?

Aufrecht has treated two hundred and ninety-four alcoholic patients between the years 1880 and 1890, and of this number has lost ten. He treated twenty-two patients with chloral and morphine, and four died; since then he has used chloral alone without any failure. He gives chloral in doses of sixty grains, according to the following formula:

R. Chloralis	3 <i>i</i>
Syripi	
Tinct. Aurantii Am. Cort.	<i>aaf</i> $\frac{3}{5}$ ss

M.

This potion, he says, is well borne by patients. It takes effect the first night, but it should be repeated the next night and the night following, so that repose may be complete. At times, in serious cases, thirty grains of chloral should be prescribed in the morning and the same dose (60 grains?) continued every evening for four or five days.

When one has to deal with all sorts of patients who at the same time are alcoholic, it is necessary to treat them actively. Aufrecht gives six fluid ounces a day of Hungarian wine, or the following potion:

R. Alcoholis (90 per cent.)	<i>f</i> $\frac{3}{2}$ ii
Syripi	<i>m</i> clx
Tinct. Aromat.	
Tinct. Amar.	<i>aa</i> $\frac{m}{xv}$
Aq. Amygdale. Amar.	<i>m</i> iii
Aq. destill.	<i>f</i> $\frac{3}{2}$ viss
Sacchari	q.s.

M.

At times, he says, this treatment does not

stop the delirium. He then prescribes forty-five grains of chloral, and has obtained very good effects from it. He says that patients with alcoholic delirium, who have no other affection, do not need alcohol.

Reduction of Hip Dislocations.

Mr. H. Herbert, Surgeon I. M. S., Acting Civil Surgeon at Aden, writes to the *British Medical Journal*, November 8, 1890: An adult Arab was admitted with an old dorsal dislocation of right hip. He had fallen from a camel quite a month before on the outer side of the limb. Thus the displacement was probably direct, and produced by forcible adduction. Hence, possibly, the failure to reduce it by the orthodox manipulations, based on the assumption of all cases being conversions of primary thyroid dislocations. Manipulations and pulleys were alternately tried for quite an hour under chloroform; but the pulleys were practically of no use, owing to the thigh band being too large. At the end of this time the limb could be abducted freely, though scarcely at all at first; and when the limb was pulled vertically upwards there was considerable movement of the head of the bone grating on the ileum. Still we quite failed with the regular movements; so, while an assistant, seizing the limb below the bent knee, pulled forcibly upwards, with the thigh fully abducted and flexed at rather more than a right angle, I, kneeling on the sound side of the patient, wrapped my arms round the limb near the hip, and, leaning my weight on the inside of the knee, so levered the head of the bone up into the acetabulum. The reduction has been permanent. The great ease with which the bone went in suggests that this method may be of somewhat general application. In old cases it probably needs free preliminary movements to break up adhesions. The extension may, of course, be made with pulleys. The essential thing is the maintenance of full abduction, while the head of the bone is pressed up into position.

Bonducine as a Febrifuge.

Bonduc is said by the *Journal de Médecine*, Oct. 12, 1890, to be the fruit of *Guilandina Bonducella* and of the *Cesalpinia Bonduc*. The oily cotyledons contain a bitter, resinous principle, which is extracted

by alcohol. The alcoholic extract is treated with chloroform, water is added to the chloroform solution, and, after agitation, the inferior layer, which contains the bitter principle, is evaporated and washed with ether and petroleum. A white powder is thus obtained, which is bitter, not acrid, soluble in alcohol, chloroform, acetone, acetic acid, fixed and volatile oils; it is sparingly soluble in ether and sulphide of carbon, and nearly insoluble in water, ether and petroleum.

This bitter principle, called bonducine, is soluble in hydrochloric acid, which it colors rose; with sulphuric acid the acid is an amaranth red, made more perceptible on the addition of a drop of chloride of iron. It is administered with success in intermittent fevers, in the dose of one and two-thirds to two and one-third grains. The powder is given in doses of fifteen or thirty grains a day.

The Quantity of Blood in the Brain.

Professor Roy and C. S. Sherrington (*Journal of Physiology*, vol. xi, p. 85), have examined the influence on the cerebral circulation of stimulation of sensory nerves, of the sympathetic, of voluntary movements, and of the action of drugs. The quantity of blood received by the brain is in proportion to the general blood pressure. They state, also, that the chemical products of cerebral metabolism contained in the lymph which bathes the walls of the arterioles of the brain, can produce variations in the caliber of the cerebral vessels, whence it follows that the brain possesses an intrinsic mechanism by whose aid the quantity of blood is made to vary locally in relation to the local variations of functional activity.—*Glasgow Medical Journal*, October, 1890.

Hydrazoic Acid.

Professor Curtius has recently discovered a body which is gaseous, and which has the formula HN_3 . It is made by the interaction of two molecules of hydrazine (N_2H_4) with one of benzoylglycollate. Benzoyl-hydrazine is one of the products, and this on heating with nitric acid is changed to the nitroso compound, which on dehydration is changed to benzoylazoinide, and saponification does the rest.—*Chemist and Druggist*, November 1, 1890.

To avert
over Confinement
and Confusion
especially VA
TEN'S O
which
performs
the nerve
refreshing
nourishing
age. One
VA H
0000A
vines ever
its growth
by its offic
DA

BA

OPI
Continu
lyn, N.
devotes
and Sel
privacy
based on
disease.

\$1,

To avoid the
well known effects of Tea
and Coffee, we con-
siderately VAN HOUTEN'S COCOA,
which is a
refreshing and
nourishing bever-
age.

One trial of
VAN HOUTEN'S
COCOA will con-
vince every one of
its great superior-
ity in strength, fla-
vor, and economy.

VAN HOUTEN'S COCOA

PURE,
SOLUBLE,
Delicious
THE STANDARD COCOA OF THE WORLD.
A SUBSTITUTE FOR TEA & COFFEE.
Easily Digested—Made Instantly.

HIGHEST AWARDS AT
PRINCIPAL EXHIBITIONS,
The Original—Take no other.

"Best & Goes Farthest—Largest Sale in the World—Once Tried, Always Used."

DANDRUFF AND BALDNESS

LASSAR'S TREATMENT *
includes Shampooing the head thoroughly for ten
minutes every day with Packer's Tar Soap.—f.f.

*Reprint from Dr. Jackson's paper—GAILLARD'S MED. JOUR., May—sent with sam-
ple of our Pine Tar Soap on receipt of 4 stamps. Address

THE PACKER MFG. CO., 100 Fulton St., New York.

HOME FOR HABITUÉS. OPIUM, CHLORAL, COCAINE.

DR. J. B. MATTISON,

Continues to receive at his residence, 314 State Street, Brooklyn, N. Y., a limited number of these habitués, to whom he devotes his *exclusive* professional attention. Patients Six, and Select. Attractive apartments, liberal cuisine, desirable privacy, cheerful society, and *personal* professional attention, based on several years' experience in the treatment of this disease. Details on application.

A GOOD PRICE WILL BE PAID FOR
the Board and Care of
a nervous lady. A physician's family preferred.

Address

DR. G. F.,

Care of Medical and Surgical Reporter, Philadelphia.
P. O. BOX 843.

CAPITAL.

\$1,000,000.

The Guarantee Trust and Safe Deposit Company,

316, 318 and 320 CHESTNUT STREET, Philadelphia,

RENTS SAFES in its ABSOLUTELY
FIRE AND BURGLAR PROOF
VAULTS.

ALLOWS INTEREST on deposits of money, acts as Registrar and Transfer Agent of Corporation Stocks, and executes Trusts of every kind under appointment of States, Courts, Corporations, or individuals, holding Trust Funds separate and apart from the assets of the Company.

COLLECTS INTEREST OR INCOME.

RECEIVES FOR SAFE KEEPING, under Guarantee, VALUABLES of every description.

Receipts for and safely keeps Wills without charge.

For further information, call at the office, or send for a circular.

MANAGEMENT.

RICHARD V. COOK, President.
GEORGE H. EARL, JR., Vice-President.
HARRY J. DELANY, Treasurer.
JOHN JAY GILROY, Secretary.
RICHARD C. WINSHIP, Trust Officer.

DIRECTORS.

Thomas Cochran.
Edward C. Knight.
Thomas MacKellar.
John J. Stadiger.
Clayton French.
W. Rotch Water.
Alfred F. tier.
J. Dickinson Sergeant.
Aaron Fries.
Charles A. Sparks.
Joseph Moore, Jr.
Richard V. Cook.
George H. Earle, Jr.



New Remedies and New Combinations.

Pil. Phenacetine et Caffeïe Citrat., W. H. S. & Co.

{ Phenacetine-Bayer, 3 grs.
Caffeïe Citrat., 1½ grs. }

Recommended in cases of Migraine and Nervous Headaches.

Pil. Phenacetine et Salol, 5 grs., W. H. S. & Co.

{ Phenacetine-Bayer, 2½ grs.
Salol, 2½ grs. }

Suggested by Dr. M. F. Price as an anti-rheumatic and analgesic. He says: "I have lately used Phenacetine largely, and have not been disappointed in its action, particularly as an analgesic. I have used it in nearly all cases of pains, especially neuralgic. I have learned to so thoroughly trust it to relieve pain that I often prescribe it without a searching inquiry into the character of the pain. A patient says, 'I have headache,' and I order Phenacetine with confidence, and always with a report of relief."

"It will be observed that in some of these cases I have combined Salol with Phenacetine. I did this on the principle of the well-known effect of Salicylic Acid in Rheumatism, but I rely on the Phenacetine for the relief of the pain, and in this way, perhaps, the cure of the disease causing it."—*Southern California Practitioner*.

Pil. Phencetine et Salol, W. H. S. & Co., ½ strength.

{ Phenacetine-Bayer, 1¼ grs.
Salol, 1¼ grs. }

Pil. Arsenici Sulphidi, 1-100 gr., W. H. S. & Co.

Dr. McLaughlin (Am. Med. Ass'n) finds Sulphide of Arsenic to be tolerated by the stomach better than arsenic in other forms. He gives it in the crude form (Realgar, As 2S2) in granules of 1-100 grain (0.0006 grammes) "ordered from Schieffelin & Co., of New York." In one patient who had suffered from psoriasis more or less for twenty years it was of great service. The eruption was on the arms, back, head, and forehead; his hands were covered with fissures, which bled when the fingers were extended; the nails were black and roughened. One granule increased to 8 granules, t. i. d., in mouth caused the fissures to heal. The face and hands were practically well. The drug was stopped for one week, when the eruption spread very rapidly. The granules were again given and improvement followed in a few days. The general health improved rapidly. In acne and eczema he has obtained satisfactory results. In every case the patient was unaware that arsenic was being administered.—*Journal Am. Med. Ass'n*.

Tablets Sulfonal et Phenacetine, 15 grs., W. H. S. & Co.

{ Sulfonal-Bayer, 10 grs.
Phenacetine-Bayer, 5 grs. }

Tablets Sulfonal et Phenacetine, 7½ grs., W. H. S. & Co.

{ Sulfonal-Bayer, 5 grs.
Phenacetine-Bayer, 2½ grs. }

For full information regarding our Soluble Pills we refer to our formula books and price-lists, which we shall be happy to furnish upon application.

W. H. Schieffelin & Co.,

170 and 172 William Street,

NEW YORK.

SUCCUS ALTERANS

(McDADE.)

SUCCUS ALTERANS is a purely vegetable compound of the preserved juices of *Stillingia Sylvatica*, *Lappa Minor*, *Phytolacca Decandra*, *Smilax Sarsaparilla*, and *Xanthoxylum Carolinianum*, as collected by Dr. GEO. W. McDADE, exclusively for ELI LILLY & CO., and endorsed by Dr. J. MARION SIMS.

SUCCUS ALTERANS continues to gain favor from its remarkable Alterative and Tonic properties, *eliminating specific poison from the blood and increasing the proportion of red corpuscles in anemic patients to a wonderful degree*; is endorsed by the medical profession, and in use by many hospitals of note. **SUCCUS ALTERANS** in venereal and cutaneous diseases is fast supplanting Mercury, the Iodides, and Arsenic; and is a certain remedy for Mercurialization, Iodism, and the dreadful effects often following the use of Arsenic in skin diseases.

SUCCUS ALTERANS is also strongly recommended for its Tonic and Alterative effects in myriad forms of scrofulous disease, and in all cases where anemia is a factor. Such patients rapidly develop a good appetite, sleep soundly, and gain flesh rapidly. Many cases are on record where patients increased ten to twenty-five pounds in weight in a few weeks.

SUCCUS ALTERANS is giving satisfactory results in treatment of *Chronic Rheumatism*, and can be used with confidence.

SUCCUS ALTERANS may be given for any length of time, without injury to the patient.

SUCCUS ALTERANS is put up in pint round amber bottles, and *never in bulk*.

Price, \$2.00 per bottle.

Send for copy of our Hand-Book of Pharmacy and Therapeutics. Useful for reference and contains much valuable information.

PIL. APHRODISIACA

(LILLY.)

Phosphorus and *Nux Vomica*, as is well known to the profession, act as powerful tonics to the nervous system, especially the spinal cord, and can be relied upon as possessing real aphrodisiac power. The Damiana used is the genuine *Turnera Aphrodisiaca*. By our process for the manufacture of Phosphorus Pills, a thorough subdivision of phosphorus in the mass is obtained, and, with a coating perfectly protecting it from oxidation, there is nothing to be desired. It is necessary that the administration of this pill be continued from three to four weeks, or until the system is thoroughly under the influence of the remedy. It is indicated in mental overwork, sexual debility, impotency. It is decidedly beneficial in cases of nocturnal emissions, the result of excess, mental apathy, or indifference, and in an enfeebled condition of the general system, with weakness or dull pain in the lumbo sacral region. In diseases of the reproductive organs of the female, and especially of the uterus, it is one of our most valuable agents, acting as a uterine tonic, and gradually removing abnormal conditions, while at the same time, it imparts tone and vigor; hence, it is of value in *Leucorrhea*, *Amenorrhea*, *Dysmenorrhea*, and to remove the tendency to repeated miscarriages.

Obtain only in original package. Oval pink pills.

One Hundred Mailed on Receipt of \$1.00.

**ELI LILLY & CO., PHARMACEUTICAL CHEMISTS,
INDIANAPOLIS, IND., U. S. A.**

SUPPLIED BY ALL DRUGGISTS.

CHARLES LENTZ & SONS,
MAKERS OF
Surgical Instruments and Orthopædic Apparatus
18 NORTH ELEVENTH STREET, PHILADELPHIA.
WE HAVE EVERYTHING IN PHYSICIANS' SUPPLIES.


Medical Batteries, Hand and Buggy Cases,
 Trusses, Elastic Hosiery.

**MICROSCOPES
 FOR
 Urinary Analysis
 A SPECIALTY.**



\$28.00 AND UPWARDS.
 Send for catalogue of Microscopes, 100 pages.

Lentz's Aseptic Catheter.



The above is a modification of Jennison's Uterine Douche, for intra-uterine injections. Any bulb or fountain syringe can be attached.

Price, with Rubber Tubing, Net, \$2.00.
 Send for descriptive Circular.

Our No. 10 Operating Case, of which so much has been said, is having large sale yet. It is the most complete set in the market for any ordinary operation. Price, \$18.75, and upward. Send for circular.

SEND FOR SPECIAL LIST OF NEW INSTRUMENTS.

ANTISEPTIC DRAINAGE TUBES.—Glass.



These Tubes have large holes, one-half inch apart, arranged alternately on opposite sides. They are carefully finished, especial care being taken to make them smooth. In addition to the drainage-holes, each tube has at one end two smaller holes for the insertion of Safety-Pin, through which it is prevented slipping into the wound.

FURNISHED IN SEVEN SIZES.
 No. 1, \$1.25 per dozen. No. 4, \$1.55 per dozen.
 No. 2, 1.25 " No. 5, 1.70 "
 No. 3, 1.40 " No. 6, 1.90 "
 No. 7, \$2.10 per dozen.

RAW CAT-GUT. I put this up in coils of 10 feet, in all different sizes, Nos. 1, 2, 3, 4 (4 is thickest). Nos. 2 and 3 are the most useful sizes. No. 1 Coil, 10 Cents; No. 2 Coil, 12 Cents; No. 3 Coil, 14 Cents; No. 4 Coil, 16 Cents. Full directions with each coil for making it absolutely aseptic.

WILLIAM SNOWDEN,
 Manufacturer, Importer and Exporter of Surgical Instruments,
 No. 121 SOUTH ELEVENTH STREET, PHILADELPHIA.

ORDER a

POCKET RECORD.

LAPACTIC PILLS, S.&D.'s.

(PILULE LAPACTICÆ.)

Aloin	1-4 gr.
Strychnine	1-60 gr.
Ext. Belladonna	1-8 gr.
Ipecac	1-16 gr.

An elegant and most efficient combination introduced by us and found in practice to possess superior advantages over similar formulæ for the relief of

HABITUAL CONSTIPATION, ATONIC DYSPEPSIA, BILIARY ENGORGEMENT AND MANY GASTRIC DISORDERS.

Since we introduced our Lapactic Pills—some four years ago—a number of manufacturers have adopted the same formula and have furnished these pills under the same name. Should Physicians fail to obtain satisfactory results from Lapactic Pills not of our make (and we have received a large number of such complaints), we shall be glad to furnish a sample of our Lapactic Pills on application. We feel confident that they will fully substantiate our claims.

Please use the term "Lapactic Pills, S. & D.'s" when prescribing these pills.
Dose : One to two pills when going to bed.

PAN-PEPTIC TABLETS, S.&D.'s.

A TONIC DIGESTIVE.

Pure Pepsin	1 gr.
Pure Pancreatin	1 gr.
Pure Caffeine	1/4 gr.
Acid Lacto-Phosphate of Calcium	
Celery	

The Pepsin and Pancreatin used in these tablets possess highest digestive power, and cannot fail to promptly start and accomplish

THE DIGESTION OF FOOD.

Whilst the Caffeine, by its stimulant tonic action on stomach and bowels, assists and quickens the normal digestion and assimilation of food, the Acid Lacto-Phosphate of Calcium contributes to the tonic action of the tablets, and aids to build up the general system, and a small quantity of that refreshing aromatic nerve stimulant, Celery, imparts a pleasant flavor and acts as an appetizer.

~~As~~ SAMPLE SENT TO PHYSICIANS WHO MENTION THIS JOURNAL.

SHARP & DOHME,

MANUFACTURERS OF

STANDARD MEDICINAL FLUID AND SOLID EXTRACTS; SOLUBLE HYPODERMIC TABLETS; SOLUBLE GELATIN AND SUGAR-COATED PILLS AND GRANULES; GRANULAR EFFERVESCENT SALTS; COMPRESSED LOZENGES AND TABLETS; FINE CHEMICALS, &c., &c.

ESTABLISHED 1860.

RABUTEAU'S DRAGEES OF IRON

Laureate of the Institute of France.—Prize in Therapeutics.
The studies made by the Physicians of the Hospitals have demonstrated that the Genuine Dragees of Iron of Rabuteau are superior to all other preparations of Iron in cases of Chlorosis, Anæmia, Leucorrhœa, Debility, Indigestion, Convalescence, Weakness of Children, and the maladies caused by the Impoverishment and Alteration of the blood after periods of fatigue, watching, and excesses of any kind.

TAKE 4 TO 6 DRAGEES DAILY.

Rabuteau's Elixir of Iron is recommended to those persons who may be unable to swallow the Dragees. Dose—*A small swallowful with meals.*

Rabuteau's Syrup of Iron is specially designed for children. Chalybeate medication, by means of Rabuteau's Iron, is the most economical and the most rational known to therapeutic.

No constipation, no diarrhea, complete assimilation.

Take only the GENUINE IRON of RABUTEAU of
CLIN & CO., Paris.

**CAPSULES
MATHEY-CAYLUS**

WITH THIN ENVELOPE OF GLUTEN.
CONTAINING COPAIBA AND ESSENCE OF SANTAL;
COPAIBA, CUBEB, AND THE ESSENCE OF SANTAL;
COPAIBA, IRON, AND THE ESSENCE OF SANTAL.

The Mathey-Caylus Capsules, of the Essences of Santal, associated with the Balms, possess an incontestable efficaciousness, and are employed with the greatest success for the rapid cure of old or recent Discharges, Gonorrhœa, Benorrhœa, Leucorrhœa, Opistho of the Neck, Urticaria, Colic, and other diseases of the bladder, together with all afflictions of the Urinary Passages."

Thanks to their thin envelope of Gluten, which is essentially assimilable, the Mathey-Caylus Capsules are digested by the most delicate persons, and never weary the stomach."

*Gentile des Hôpitaux de Paris,
CLIN & CO., Paris,
AND OF ALL DRUGGISTS.*

**SOLUTION OF
THE SALICYLATE OF SODA
OF DOCTOR CLIN.**

Laureate of the Paris Faculty of Medicine
(MONTYON PRIZE).

Dr. Clin's Solution, always identical in its composition, and of an agreeable taste, permits the easy administration of pure Salicylate of Soda, and the variation of the dose in accordance with the indications presented.

"The Salicylate of Soda used by Clin is of perfect purity, and is prepared with the greatest care; it is a medicament in which we may have every confidence."

—Paris Society of Medicine, Meeting of Feb. 8th, 1879.
Clin's Solution, very exactly mixed, contains: 8 grammes of Salicylate of Soda per tablespoonful, 60 centigrammes of Salicylate of Soda per teaspoonful.

PARIS-CLIN & CO.—PARIS
AND BY ALL DRUGGISTS.

**NEURALGIAS
PILLS OF DR. MOUSSETTE.**

The Moussette Pills of aconitine and quinum, calm or cure Gastralgia, Hemisconia, Headache, Sciatic, and the most obstinate Neuralgias.

"The sedative action exerted by the Moussette Pills upon the apparatus of the sanguineous circulation by the intercession of the vaso-motor nerves, indicates their use in neuralgias of the trigeminal nerve, (fifth pair), congestive neuralgia, and painful and inflammatory Rheumatism affections."

"Aconitine produces marvelous effects in the treatment of facial neuralgias when they are not symptomatic of intracranial tumor."—*Society of Biology of Paris, Meeting of the 28th February, 1880.*

Dose—Take 3 to 6 pills during the twenty-four hours.

**ACCEPT ONLY THE GENUINE MOUSSETTE PILLS OF
CLIN & CO.—PARIS.**

GRAND NATIONAL PRIZE OF 16,600 FRANCS,**QUINA-LAROCHE**

This meritorious Elixir, QUINA-LAROCHE, is prepared from the three Cinchonas; it is an agreeable and doubtless highly efficacious remedy.

—The Lancet.

**VINOUS ELIXIR,
A STIMULATING
RESTORATIVE
AND
ANTI-FEBRILE TONIC.
FAR SUPERIOR TO ALL ORDINARY CINCHONA WINES.**

QUINA-LAROCHE
under the form of a vinous Elixir contains the totality of the numerous principles of the three Cinchona barks. It is a preparation which renders great services to Therapeutics.—Extract of the *Gazette des Hôpitaux*, Paris.

LAROCHE'S QUINA, tried by the Academy of Medicine of Paris, is a specially prepared COMPOUND EXTRACT of Quinquina, a careful analysis, confirmed by experience, has shown that most of the wines and syrups hitherto used have not contained all the properties of this precious bark, of these some, although beneficial, are altogether lost, while many preparations contain but half the properties of the bark in varying proportions.

MR. LAROCHE, by his peculiar method, has succeeded in extracting all the active principles of the quinquina, combining these with Cinchona Wine forming an Elixir free from the disagreeable bitterness of other similar preparations. Practitioners have found triple advantages in the use of this Elixir; it is a strong tonic, is easily administered, and perfectly harmless, being free from the unpleasant effects of Quinine.

THE FERRUGINOUS QUINA-LAROCHE is the invigorating tonic par excellence, having the advantage of being easily assimilated by the gastric juice; without, in any way, deranging the action of the digestive organs, proving itself to be a most efficacious remedy in cases of impoverishment of the blood, ANÆMIA, CHLOROSIS, INTESTINAL HEMORRHAGE, CONSTIPATION, EXHAUSTION, ETC., ETC.

PARIS—22 RUE DROUOT—PARIS.

E. FOUGERA & CO., New York,

Sole Agents for the United States for the above Preparations.

TERTIARY SYPHILIS.

"Your preparation, **IODIA**, is an excellent alternative, and is the most agreeable of any preparation of Iod. Potas., as it does not cause the cramps in stomach, which are often intolerable.

"I have employed **IODIA** in two cases of Tertiary Syphilis, in which, after five weeks' treatment, the manifestations disappeared.

"In a case of Syphilitic Eczema it was also successful.
—DR. SAM PETITEAU, Garches, France, Nov. 10, 1890.

BATTLE & CO., CHEMISTS' CORPORATION,

St. Louis, Mo., U. S. A.

BRANCHES:

- 76 New Bond Street, London, W.
- 5 Rue de la Paix, Paris.
- 9 and 10 Dalhousie Square, Calcutta.
- 80 Montague-de-la-Cour, Brussels.
- 28 Nieuwe Hoogstraat, Amsterdam.

HELP!!

WE WANT TO ASK all the FRIENDS of the MEDICAL AND SURGICAL REPORTER to
HELP to make it the strongest **AMERICA**, or anywhere,
 Medical Journal in for that matter.

It now has subscribers in every **STATE AND TERRITORY**, in *Canada, Mexico, South America, Europe, Asia, and AFRICA.* But it could have even more. *We might employ Convassers* to go around and solicit subscriptions, but we would prefer to have ITS FRIENDS recommend it to their friends. You KNOW WHAT IT IS and can tell.

It aims to be

INTERESTING!
INSTRUCTIVE!
WIDE-AWAKE!
FEARLESS!

If it is **NOT**; then
 don't recommend it.

And not *too heavy* for a *busy Doctor* to enjoy.

We don't **ASK ANYBODY** to do { *something*
for nothing, } so we propose to offer to our subscribers the following **PREMIUM FOR NEW SUBSCRIBERS.** To any paid-up subscriber sending us, before January 1, 1892, the name of one new subscriber (with \$5.00), we will send as

A PREMIUM

any publication or instrument in our own list, or that of any publisher or instrument maker, to the value of **TWO DOLLARS.** OR: any subscriber sending a *new name and eight dollars*, will be credited with a *year's subscription* on his own account, and the REPORTER will be sent to the new address until **January 1, 1892.**

SEND CHECK OR MONEY ORDER TO

MEDICAL AND SURGICAL REPORTER,

P. O. Box 843.

PHILADELPHIA.

SPECIAL OFFER

**TO SUBSCRIBERS TO THE REPORTER WE MAKE
THE FOLLOWING OFFER:**

FOR TEN DOLLARS WE WILL SEND

The REPORTER for one year,	price alone, \$5.00
Model Ledger,	.50
Accidents and Emergencies,	.75
Pocket Record,	1.25
Total,	\$12.00

FOR NINE DOLLARS WE WILL SEND

The REPORTER for one year,	price alone, \$5.00
Model Ledger,	5.00
Pocket Record,	<u>1.25</u>
Total,	\$11.25

FOR EIGHT DOLLARS WE WILL SEND

The REPORTER for one year, Model Ledger, Accidents and Emergencies,	price alone, \$5.00 5.00 .75
Total,	\$10.75

FOR SIX DOLLARS WE WILL SEND

The REPORTER for one year, Pocket Record, Accidents and Emergencies,	price alone, \$5.00 1.25 .75
Total,	<u>\$7.00</u>

Send Check or Money Order to

MEDICAL AND SURGICAL REPORTER,

P. O. BOX 843.

PHILADELPHIA.

BOOKS FOR OUR SUBSCRIBERS

BY SPECIAL ARRANGEMENT with the publishers we are able at this time to offer the following first-class and very valuable books (last editions) at a very low price, when taken in connection with paid-up subscriptions. Subscribers will see that we have put in this list thoroughly good books. See Adv. Page xi.

FOR **\$10.00**

REPORTER for one year,	\$5.00
HEATH'S DICTIONARY OF PRACTICAL SURGERY. Cloth,	\$7.50

FOR **\$9.00**

REPORTER for one year,	\$5.00
LEIDY'S HUMAN ANATOMY. By Joseph Leidy, M. D. Cloth,	\$6.00

FOR **\$8.50**

REPORTER for one year,	\$5.00
and any one of the following books—	
1.—BYFORD'S DISEASES OF WOMEN. Cloth,	\$5.00
2.—FOWLER'S DICTIONARY OF PRACTICAL MEDICINE.	\$5.00
3.—JACOBSON'S OPERATIONS OF SURGERY,	\$5.00
4.—CAZEAX & TARNIER'S MIDWIFERY. With Appendix by Munde,	\$5.00
5.—THOMAS'S MEDICAL DICTIONARY,	\$5.00

These works are all in cloth binding, but can be supplied in leather binding for \$1.00 additional.

FOR **\$6.00**

we will send the **REPORTER** for one year, **\$5.00**
and any two of the following books:

1.—THE NURSING AND CARE OF THE NERVOUS AND THE INSANE. By Chas. K. Mills, M. D.,	\$1.00
2.—MATERNITY; INFANCY; CHILDHOOD. By John M. Keating, M. D.,	\$1.00
3.—OUTLINES FOR THE MANAGEMENT OF DIET; or, The Regulation of Food to the Requirements of Health and the Treatment of Disease. By E. T. Bruen, M.D.,	\$1.00
4.—FEVER NURSING. Designed for the use of professional and other Nurses. By J. C. Wilson, A. M., M. D.,	\$1.00
5.—DISEASES AND INJURIES OF THE EAR: Their Prevention and Cure. By Chas. H. Burnett, A. M., M. D.,	\$1.00

Or, For **\$6.00**, any one of the above Nursing Books and **THOMSON'S (Sir Henry) SURGERY OF THE URINARY ORGANS.** Some important points connected with the Surgery of the Urinary Organs. Illustrated. Cloth, **\$1.25**

Books for our Subscribers.

WHAT IS THOUGHT OF THEM.

HEATH'S DICTIONARY OF PRACTICAL SURGERY.

C. B. Porter, M. D., Boston, Mass., says: "A most excellent book for the library of the surgeon, and especially for the country practitioner; as a book of reference it is so concise and at the same time so complete." *N. Senn, M. D., College of Physicians and Surgeons, Milwaukee*, says: "As a means of ready reference for the student and busy practitioner this book stands unexcelled."

Prof. E. H. Gregory, St. Louis Medical College, St. Louis, says: "I have examined the book, and find it a most valuable addition to my library. I am sure it will meet the favor of the surgical profession."

LEIDY'S HUMAN ANATOMY.

The Lancet-Clinic, Cincinnati, O., says: "Professor Leidy again presents to medical practitioners and students a treatise on human anatomy that at once commands the attention and admiration of all who are at all familiar with the subject. Most of the plates are original, and in common with the text are very beautiful to behold. The work is a complete illustration of the method of teaching anatomy adopted by a leading scientist, and one of America's best known professors of anatomy. This is one of the books that should be in the library of every practitioner of medicine and surgery."

The Medical Advance, Ann Arbor, Mich., says: "The student can master and retain a practical knowledge of anatomy in a shorter time and with less hard work from this text-book than from any other work extant, and it has been our privilege to teach anatomy for several years."

BYFORD'S DISEASES OF WOMEN.

The Boston Medical and Surgical Journal says: "It is a satisfaction to observe the discussion of certain topics which are often neglected by writers, notably of the sympathetic symptoms of organs apparently disconnected with the uterus, for which he will receive the thanks of the younger members of the profession. Another topic very widely treated is that of Constipation in its relation to uterine diseases. All the other topics which should have a place in a work of this kind have been fully and properly treated."

The Cincinnati Lancet-Clinic says: "The book is eminently practical and instructive. Its suggestions for treatment are well up with the times, mostly judicious, conservative, and valuable, and the whole amply repays for thorough perusal."

The St. Louis Courier of Medicine says: "The work of a representative Western practitioner and professor, and is a thoroughly practical work."

The Virginia Medical Monthly says: "We take pleasure in recommending it to every student and physician desirous of obtaining a standard work on Gynecology."

The Chicago Medical Journal and Examiner says: "The chapter on puberty should be carefully studied by every physician, teacher, and mother in the land. . . . Its style is clear, but concise, and always practical, and shows the author to be, as we all know, an experienced teacher. The work as a whole is most valuable, both to the student and practitioner."

Gaillard's Medical Journal says: "It is replete with good judgment, broad and practical views. . . . The illustrations are excellent."

FOWLER'S DICTIONARY OF PRACTICAL MEDICINE.

The Medical News says: "For those who do not wish to purchase the much larger book of Quain, or who wish comparatively recent information concerning medical subjects in a somewhat narrow space, the book will certainly prove of value, and we heartily recommend it as one which will give a full return for the money invested in the purchase."

The London Lancet says: "What the writers know they tell simply and plainly, and generally manage to put it in a short compass, and, what is more, the treatment that is recommended is clearly and sufficiently set forth. . . . We can heartily congratulate the editor and his colleagues on having admirably fulfilled their purpose, viz., to produce a first-rate dictionary of medicine, at once scientific and practical, which well deserve the success we have no doubt it will meet with."

JACOBSON'S OPERATIONS OF SURGERY.

The Medical and Surgical Reporter says: "This handsome book is one of the most admirable works on operative surgery which we have seen for a long time. The author has brought to its preparation a large observation and experience, and a very thorough acquaintance with the methods of other surgeons. His book is comprehensive and his directions are clear and reliable. It is gratifying to an American reader to find that full justice is done to the work of American surgeons by Mr. Jacobson, and that his book bears no marks of the distance which stretches between his land and ours. This is as it should be, and is no less an element in the excellence of the technical worth of his book than it is an evidence of the cosmopolitan character of good scientific work."

CAZEAUX & TARNIER'S MIDWIFERY. With Appendix by MUNDÉ.

The Memphis Medical Monthly says: "Cazeaux's Obstetrics was authority during our student life, and has been generally recognized as authority from the publication of the first edition to the present. We cannot do more than to fully endorse the following from the pen of T. Gallard Thomas, M. D.: 'I have examined the last edition of Cazeaux & Tarnier's Theory and Practice of Obstetrics, and pronounce it practical, and just what is needed by every practitioner. I highly recommend the work. This work should be prominent in every library.'

Dr. Wm. L. Lush says: "Cazeaux's Midwifery will always remain an inexhaustible mine of information, and I rejoice to see it once more accessible to the profession in a new and attractive dress."

Dr. Wm. Goodell says: "I deem Cazeaux & Tarnier's Obstetrics the best yet published."

The Denver Medical Times says: "It is needless to say anything in praise of this work. Cazeaux's Obstetrics is a classical work, and Tarnier is one of the best known French writers on Obstetrics. The whole has been brought up to date by the appendix of Prof. Mundé, and altogether it makes a very complete and practical treatise on the subject, unsurpassed if not unequalled by any work with which we are acquainted. Especially valuable are the signs of delivery in the dead body and of live and still-births. The chapter on antiseptics and on puerperal septicemia are also worthy of mention."

THOMAS'S MEDICAL DICTIONARY.

Prof. A. F. Patton, College of Physicians and Surgeons, Boston, Mass., says: "It is just the book for a medical or any other student, and it should be in the office of every physician. This dictionary supplies a place that has never been filled. I have looked it through and find all the new words that I have sought."

The St. Joseph Medical Herald says: "No better testimonial to the value of the work can be given than the following from Dr. J. Gibbon Hunt, the distinguished microscopist: 'To me the work is invaluable. I am astonished at its fullness of all recent words which the modern advance of medical science has introduced. Of course, I cannot conceive the learning and great labor which could edit such a complete, thorough, and admirable volume.' We can add nothing further save that the book should be in the hands of every physician and every student in the land."

DISEASES IN WHICH Oxygen and Nitrogen Monoxide HAVE BEEN EMPLOYED.

(See "THERAPEUTIC USES OF OXYGEN AND NITROGEN MONOXIDE," which can be obtained upon application.)

Anæmia,	Dyspnoea,	Nephritis,
Asphyxia,	Emphysema, Pulmonary,	Nervous Affections,
Asthena,	Epilepsy,	Nervous Prostration,
Asthma,	Formication,	Neuralgia,
Atonic Conditions,	Headache,	Paralysis,
Bright's Disease,	Hemorrhage, Pulmonary,	Phtisis,
Bronchitis,	Hypochondriasis,	Pleuritic Adhesions,
Catarrh,	Hysteria,	Pleurisy,
Croup,	Indigestion,	Pleuro-Pneumonia,
Cyanosis,	Insomnia,	Pneumonia,
Diabetes,	Laryngitis,	Rheumatism.
Diarrhoea,	Lithiasis,	Scarlet Fever,
Diphtheria,	Melancholia,	Tuberculosis,
Dyspepsia,	Menstrual Irregularities,	Uræmia.

ABOUT GAS TREATMENT.

We make and sell oxygen and nitrogen monoxide for therapeutic use, and we guarantee them pure. They are put up in compact form. (A cylinder containing 100 gallons of nitrogen monoxide or 40 gallons of oxygen measures 12 inches in length, has a diameter of 3½ inches, and weighs 10½ lbs. A cylinder containing 450 gallons of nitrogen monoxide or 100 gallons of oxygen measures 25 inches in length, has a diameter of 4½ inches, and weighs 34 lbs.)

INSOMNIA. Dr. Allan McLane Hamilton states that nitrogen monoxide, N_2O (nitrous oxide of the old nomenclature) has no equal in the treatment of this difficulty.

MELANCHOLIA. A short course of nitrogen monoxide is said to change the face of nature for such patients.

ANÆMIA. Where iron is not tolerated or proves inefficient, the addition of oxygen or a combination of oxygen and nitrogen monoxide has proven very beneficial.

For those who should, but cannot, go from home for rest or a change of air, or for those who, from any cause fail to get air enough into their lungs, the inhalation of one or other of these gases, or both in combination, promises great benefit. The testimony is that in cases of

ASTHMA, more than fifty per cent, of the cases yield to oxygen treatment, others are very greatly relieved, and a very small per cent, are not improved.

INDIGESTION and **CONSTIPATION** are said to be greatly relieved and very often conquered by continued treatment.

A highly esteemed New York physician has used more than twenty thousand gallons of nitrogen monoxide in his private practice. The letters received from his patients, largely from those who were afflicted with nervous disorders, are enthusiastic in their testimony as to the benefit received.

Regular practitioners who are using gas treatment testify that when *pure* oxygen and *pure* nitrogen monoxide, or a mixture of these, is used, no therapeutic agents are more uniformly successful when intelligently prescribed.

For descriptive circulars, and term for gas outfits, address

The S. S. White Dental Manufacturing Company,

AT EITHER OF THE BELOW NAMED PLACES:

Twelfth and Chestnut Sts., Philadelphia, Pa. 160 Tremont St., Boston, Mass.
 767 and 769 Broadway (cor. 9th St.), New York, N. Y. 151 and 153 Wabash Ave., Chicago, Ill.
 1260 and 1262 Broadway (cor. 32d St.), New York, N. Y. 444 Fulton St., Brooklyn, N. Y.

**THE BEST ANTISEPTIC
FOR BOTH INTERNAL AND EXTERNAL USE.**

**ANTISEPTIC,
PROPHYLACTIC,
DEODORANT.**

LISTERINE

**NON-TOXIC,
NON-IRRITANT,
NON-ESCHAROTIC.**

FORMULA—Listerine is the essential antiseptic constituent of Thyme, Eucalyptus, Baptisia, Gaultheria and Mentha Arvensis, in combination. Each fluid drachm also contains two grains of refined and purified Benzo-boracic Acid.

DOSE—Internally: One teaspoonful three or more times a day (as indicated) either full strength, or diluted, as necessary for varied conditions.

LISTERINE is a well-proven antiseptic agent—an antizymotic—especially adapted to internal use, and to make and maintain surgical cleanliness—asepsis—in the treatment of all parts of the human body, whether by spray, irrigation, atomization, or simple local application, and therefore characterized by its particular adaptability to the field of

PREVENTIVE MEDICINE—INDIVIDUAL PROPHYLAXIS.

Diseases of the Uric Acid Diathesis.

LAMBERT'S

LITHIATED HYDRANGEA

KIDNEY ALTERATIVE—ANTI-LITHIC.

FORMULA—Each fluid drachm of "Lithiated Hydrangea" represents thirty grains of FRESH HYDRANGEA and three grains of CHEMICALLY PURE Benzo-Salicylate of Lithia. Prepared by our improved process of osmosis, it is INvariably of DEFINITE and UNIFORM therapeutic strength, and hence can be depended upon in clinical practice.

DOSE—One or two teaspoonfuls four times a day (preferably between meals).

Urinary Calculus, Gout, Rheumatism, Bright's Disease, Diabetes, Cystitis, Hematuria Albuminuria, and Vesical irritations generally.

We have much valuable literature upon **GENERAL ANTISEPTIC TREATMENT,** **LITHIUM, DIABETES, CYSTITIS, ETC.** To forward to Physicians upon request:

LAMBERT PHARMACAL CO., ST. LOUIS, MO.

FRENCH'S FLUID EXTRACTS

Are prepared from carefully selected crude drugs, ground in mills on our own premises. The most scientific and recent methods are used. Careful consideration is given as to the best menstruum for each drug. Each fluid ounce of our fluid extracts represents one troy ounce of the drug, minim for grain. Strength, dose, and formula given on the label.

FRENCH'S COMPRESSED MEDICINAL LOZENGES.

OFFICIAL LOZENGES OF THE U. S. PHARMACOPEIA, 1880

FRENCH'S COMPRESSED LOZENGES OF THE LONDON HOSPITAL FOR DISEASES OF THE THROAT.

FORMULÆ SUGGESTED BY DR. MORELL MACKENZIE.

(Made with black and red currant paste.)

FRENCH'S COMPRESSED UNOFFICIAL LOZENGES.

FRENCH'S COMPRESSED MEDICATED LOZENGES.

(SPECIAL FORMULÆ.)

Our laboratory is in charge of Professor F. G. RYAN, of the Philadelphia College of Pharmacy, who has able assistants.

Send for our new price-list with formulæ for Compressed Pills, Lozenges, Hypodermic Tablets, Tablet Triturates, etc.

FRENCH, RICHARDS & CO., PHILADELPHIA.

A MENSTRUUM.

HORSFORD'S ACID PHOSPHATE.

THIS preparation has been found especially serviceable as a menstruum for the administration of such alkaloids as morphine, quinine, and other organic bases which are usually exhibited in acid combination.

The admixture with pepsin has been introduced with advantage when indicated.

The Acid Phosphate does not disarrange the stomach, but, on the contrary, promotes in a marked degree the process of digestion.

Dr. R. S. MILES, Glencoe, Minn., says: "I use it in a great many cases as a menstruum for quinine, when an acid is necessary."

Send for descriptive circular. Physicians who wish to test it will be furnished a bottle on application, without expense, except express charges.

Prepared under the direction of Prof. E. N. HORSFORD, by the

Rumford Chemical Works, Providence, R. I.

BEWARE OF SUBSTITUTES AND IMITATIONS.

CAUTION:—Be sure the word "Horsford's" is printed on the label. All others are spurious. Never sold in bulk.

FRELIGH'S TABLETS,

(Cough and Constituent),

FOR THE PREVENTION AND CURE OF PULMONARY PHthisis. FORMULÆ.

Cough Tablets.

EACH TABLET CONTAINS.

Morph. Sulph. ($\frac{1}{10}$ gr.), Atropia Sulph. ($\frac{1}{10}$ gr.), Codex ($\frac{1}{10}$ gr.), Antimony Tart. ($\frac{1}{10}$ gr.), Ipecac, Aconite, Pusatilla, Dulcamara, Causticum, Graphite, Rhus-tox, and Lachesis, fractionally so arranged as to accomplish every indication in any form of cough.

Constituent Tablets.

EACH TABLET CONTAINS.

Arsenicum ($\frac{1}{10}$ gr.), Precipitate Carb. of Iron, Phos. Lime, Carb. Lime, Silica, and the other ultimate constituents, according to physiological chemistry (normally) in the human organism, together with Caracas. Cocoa and Sugar.

PRICE, THREE DOLLARS PER DOUBLE BOX.

Containing sufficient Tablets of each kind to last from one to three months according to the condition of the patient.

SPECIAL OFFER.

While the above formulas have been in use, in private practice, over 30 years, and we could give testimonials from well-known clergymen, lawyers and business men, we prefer to leave them to the unbiased judgment of the profession with the following offer: On receipt of 50 cents, and card, letter-head, bill-head, or other proof that the applicant is a physician in active practice, we will send, delivered, charged prepaid, one of the regular (double) boxes (retail price, Three Dollars), containing sufficient of each kind of Tablets to test them three months (in the majority of cases) in some one case. Card, letter-head, or some proof that the applicant is a physician in active practice, must accompany each application. Pamphlet, with full particulars, price-list, etc., on request.

As we furnish no samples through the trade, wholesale or retail, for samples, directions, price-list, etc., address,

I. O. WOODRUFF & CO.,

MANUFACTURERS OF PHYSICIANS' SPECIALTIES,

88 Maiden Lane,

New York City.

"Familiar in millions of mouths as any household word."—
The Times.

Apollinaris

"THE QUEEN OF TABLE WATERS."

"In 1873 the number of bottles filled and exported was a little under 2,000,000. Last year the number was nearly 16,000,000."

"Filling went on at the rate of 90,000 a day."

"These figures are gigantic."

THE TIMES, LONDON, 20th Sept., 1890.


THE BEST NATURAL APERIENT.

THE APOLLINARIS COMPANY, LIMITED, London, beg to announce that, as numerous APERIENT WATERS are offered to the public under names of which the word "Hunyadi" forms part, they have now adopted an additional Label comprising their Registered Trade Mark of selection, which consists of

A RED DIAMOND.

This Label will henceforth also serve to distinguish the HUNGARIAN APERIENT WATER SOLD BY THE COMPANY from all other Aperient Waters.

DEMAND THE



DIAMOND MARK.

And insist upon receiving the HUNGARIAN APERIENT WATER of the APOLLINARIS COMPANY, LIMITED, London.

BOUDAULT'S PEPSINE

The only Pepsine used in the Hospitals of Paris for the last Thirty Years.

Unlike the various substitutes which, in most cases, are but unscientific or incompatible compounds, forced upon the Medical Profession as aids to digestion by extensive advertising, but which, when submitted to the proper tests, are found to be useless as digestive agents, Pepsine is constantly gaining in the esteem of the careful practitioner.

Since the introduction of Pepsine by Boudault and Corvisart in 1854, the original BOUDAULT'S PEPSINE HAS BEEN AT ALL TIMES CONSIDERED THE BEST, as is attested by the awards it has received at the Expositions of 1867, 1868, 1871 1873, in 1876 at the Centennial Exposition of Philadelphia, and in 1878 at the Paris Exposition.

The most reliable test, carefully applied, will satisfy everyone that BOUDAULT'S PEPSINE HAS A MUCH HIGHER DIGESTIVE POWER than the best Pepsines now before the Profession, and is therefore especially worthy of their attention.

BOUDAULT'S PEPSINE is prepared in the form of Pepsine Acid and Pepsine Neutral. It is sold in bottles of one ounce, with a measure containing exactly five grains; also in bottles of four, eight, and sixteen ounces for dispensing.

BOUDAULT'S WINE OF PEPSINE

FORMULA OF DR. CORVISART.

The taste of Pepsine being perfectly disguised in this Wine, it may be recommended to persons who have difficulty in taking Pepsine in the form of Powder. This Wine is tested so that a tablespoonful of it is equal in digestive power to ten grains of Boudault's Pepsine in powder. Sold only in bottles of eight ounces.

TANRET'S PELLETIERINE

For the Treatment of Tape-Worm (Tania Solium).

This New Tannifuge, the Active Alkaloid of Pomegranate Bark, has of late come into extensive use in France for the treatment of Tape-Worm (Tania Solium). The results of numerous experiments with it at the Marine Hospitals of Toulon, St. Mandrier, etc., and in the Hospitals of Paris, St. Antoine, La Charité, Necker Beaujon, etc., have all been most satisfactory. Doctor Dujardin Beaumets, Member of the Academy of Medicine, and Professor Laboulbene, in their report to the Society of Therapeutics, have given it their unqualified approval, after the most searching experiments. This preparation is pleasant to administer, and, if certain preliminaries are observed, success will be insured.

Sold only in bottles containing one dose.

DUCRO'S ALIMENTARY ELIXIR

A Combination uniting the properties of Alcoholic Stimulants and Raw Meat.

This preparation, which has been used with great success in the hospitals of Paris since 1868, is adapted to the treatment of all diseases requiring administration, in a small volume, of a tonic able to stimulate and support the vital forces, as Pulmonary Fibrosis, Depression and Nervous Debility, Adynamia, Malariaous Cachexia, etc.

Prepared by EMILE DURIEZ & CO., Successors to DUCRO & CIE, Paris.

KIRKWOOD'S INHALER

This is the only complete, reliable, and effective inhaler in use, arranged for the direct application of Muriate of Ammonium and other remedial agents in the state of vapor to the diseased parts of the air-passages in the treatment of catarrh and diseases of the throat and lungs. No heat or warm liquids required in its use.

It is entirely different from the various frail, cheap instruments that have been introduced.

KIRKWOOD'S INHALER is accompanied by testimonials of the highest professional character, together with carefully prepared formulas for use.

RETAIL PRICE, COMPLETE, \$2.50.

~~AD*~~ A liberal discount allowed to the trade and profession. For descriptive pamphlet or other information address

E. FOUGERA & CO., 30 North William St., New York.

Sole Agents for the above Preparations.

icularly t
er throat,
Hydrogen
in destroy
may be en
“In
the full st
irritation
standard 1

“An
mined effe
with an a
prolonged
be follow

CAU
only in %.

A book c
medical ap
HYDROGEN
profession
application.
SOLD BY

Constipation.

Pavara Pills are prepared especially for the treatment of Constipation. They are tasteless, harmless, and produce natural movements of the bowels, without pain, nausea, or any inconvenience whatever.

Physicians are requested to test these pills in those cases which they think to be incurable, also, in cases of constipation during pregnancy. Thousands of physicians now use and prescribe them.

Physicians will see by the formula (on the outside label) that each medicinal ingredient is absolutely harmless.

A large bottle, for trial, will be mailed free to any practising physician.

Prepared at **TOPFLY'S BOTANIC LABORATORY**,
New York. Depot, 115 Fulton Street.

Please mention this Journal.

CH. MARCHAND'S PEROXIDE OF HYDROGEN

(MEDICINAL) H_2O_2 (ABSOLUTELY HARMLESS.)

Is rapidly growing in favor with the medical profession. It is the most powerful antiseptic known, almost tasteless, and odorless. Can be taken internally or applied externally with perfect safety. Its curative properties are positive, and its strength and purity can always be relied upon. This remedy is not a nostrum.

A REMEDY FOR DIPHTHERIA; CROUP; SORE THROAT,
AND ALL INFLAMMATORY DISEASES OF THE THROAT.

OPINION OF THE PROFESSION.

Dr. Geo. B. Hope, Surgeon Metropolitan Throat Hospital, Professor Diseases of Throat, University of Vermont, writes in an article headed, "Some Clinical Features of Diphtheria, and the treatment by Peroxide of Hydrogen" (*N.Y. Medical Record*, October 13, 1888). Extract:

"On account of their poisonous or irritant nature the active germicides have a utility limited particularly to surface or open wound applications, and their free use in reaching diphtheritic formations in the mouth or throat, particularly in children, is, unfortunately, not within the range of systematic treatment. In Peroxide of Hydrogen, however, it is confidently believed will be found, if not a specific, at least the most efficient topical agent in destroying the contagious element and limiting the spread of its formation, and at the same time a remedy which may be employed in the most thorough manner without dread of producing any vicious constitutional effect."

"Inasmuch as the efficacy depends upon the ozonized oxygen in solution, it has seemed desirable to rely on the full strength of the officinal preparation of fifteen volumes, especially when used in the fauces, where any slight irritation from its acidity is not apparent. In all the cases treated (at the Metropolitan Throat Hospital), a fresh, standard Marchand preparation of fifteen volumes was that on which the experience of the writer has been based."

"An equally important element is in making the application in such a manner as to produce the most determined effect on the diseased tissues with as little local disturbance as possible. . . . A steady, coarse spray, with an air pressure of twenty pounds or more, will, in a few moments' time, produce a more positive action than prolonged efforts to reach the fauces by means of cotton applicators. . . . How frequently the treatment is to be followed up depends to a considerable extent on the density, as well as the area, of the surface involved. . . ."

CAUTION.—By specifying in your prescriptions "Ch. Marchand's Peroxide of Hydrogen (Medicinal)," which is sold only in $\frac{1}{4}$ -lb., $\frac{1}{2}$ -lb., and 1-lb. bottles, bearing my label and signature, you will never be imposed upon. Never sold in bulk.

PREPARED ONLY BY

A book containing full explanations concerning the therapeutic applications of both CH. MARCHAND'S PEROXIDE OF HYDROGEN (Medicinal) and GLYCOSIDE, with opinions of the profession, will be mailed to physicians free of charge on application.

Mention this publication.

SOLD BY LEADING DRUGGISTS.

Laboratory, 10 West Fourth Street, New York.

Charles Marchand

Chemist and Graduate
of the "Ecole Centrale
des Arts et Manufactures de Paris"
(France).

A CASE WITH A LESSON IN IT.

We have frequently referred to the necessity of keeping the pregnant woman supplied with those chemical substances which she must give to the child in the formation of its little frame, and the dire effects which may result from a deficiency of those substances. The headaches, neuralgias, spinal tenderness, and some of the many disorders of digestion, and even mental failure, may be all prevented or relieved by supplying to the mother the chemical food her system demands.

We have noticed a few cases where the most serious results have occurred to the mother by such a deficiency, one of which is well worth relating. Mrs. M., aged 30, pregnant with her third child, suffered from the second to the fifth month of her pregnant term with gradually failing strength and health, and nervous irritability. She was very restless nights for three or four days, the peculiar restlessness of complete nervous exhaustion, and was irrational for more than a week.

At this time I advised that she have a teaspoonful of Murdock's Liquid Food every hour in milk, and a little wine every four hours. The third night she had fifteen drops of hydrobromic acid dil. every three or four hours, which was continued for a week. The liquid food after a week every two hours, and all other medicines were discontinued. Her bowels were moved with a mild laxative at first, and afterward became regular. Her tongue cleaned quickly and her skin assumed a more natural appearance.

At this writing, less than three weeks from our first visit, the patient is able to sit up an hour at a time, is quite rational and cheerful, has a good appetite, the bowels are quite regular, there is no headache, and but little spinal tenderness. She sleeps reasonably well if fed frequently. She is allowed to suit her taste in her diet. The child is quite active, and the mother is progressing so nicely that it is hoped that she will pass safely through her confinement, which is nearly at hand. She will be kept on the liquid food and the phosphates until her confinement after which, the condition will suggest the treatment. This is the most marked case of a starved nervous system we have ever seen, and the benefit derived from the treatment

proves the theory as to the cause of the trouble.

Apropos with the above, comes the question of the selection of nutritious foods for those many cases which demand strong and immediate nourishment. The food used in the above case has produced marvelous results for us in some of those extreme cases in which we have used it.

In one case of nervous exhaustion the insomnia was absolutely painful. When the exhaustion forced the patient to sleep there was an immediate sense of falling or approaching death, or there were hallucinations which were appalling, rendering the nights unbearable. One teaspoonful of Murdock's Liquid Food so nourished the system the first night it was given that the patient sank into a quiet, refreshing sleep which lasted, with a single awakening, until morning, and the improvement the patient obtained was most striking.

One case of mitral regurgitation with leucocythemia and nervous exhaustion improved more rapidly on this food than on all other measures combined. In cases of nervous exhaustion and brain tire, its effects are most marked, as we have experienced many times in the past five years in the frequent personal use of the agent. It contains the principles of pure, healthy blood and nerve tissue ready for immediate assimilation. Fourteen per cent. of the entire quantity is uncoagulated albumen, as we have demonstrated by actual analysis.

We have tided babies over critical periods often with a few drops in its usual diet. We have given it to dipsomaniacs with or without non-alcoholic stimulants with superb results. The sole objection our patients have to it is its odor. Its taste is not disagreeable; it has the odor and appearance of blood. All objections are soon forgotten, however, in the results of its action upon the nutrition of the system, and the patient soon learns to like it, and often craves for it, drinking it from the bottle, or a glass readily, like any beverage. It ought to be the food par-excellence for rapid appropriation in all conditions of exhaustion.

P. S.—Phosphates act well with the Liquid Food.

Editorial of the *Chicago Medical Times*, June, 1890.

**MESSRS. SNOW, CHURCH & CO.,
LAW AND COLLECTION OFFICES,
608 & 610 Chestnut Street,
PHILADELPHIA.**

GENTLEMEN:—Your management of the collections of the MEDICAL AND SURGICAL REPORTER during the past year has been so admirable and so much more successful than that of any agency with which I have dealt heretofore, that I renew my contract with you for the coming year with great pleasure.

The Publisher of the Medical and Surgical Reporter.
Philadelphia,
September 25, 1890.

GENUINE BLAUD'S PILLS.

These pills, which have been inserted in the new French Pharmacopœia, have been employed with the greatest success for more than fifty years by most French and foreign physicians, to cure anaemia, chlorosis, and all chlorotic affections in which iron is indicated.

Here is the opinion of men most eminent in medical science who have employed them:

"For thirty-five years, in which I have practised medicine, I have recognized the incontestable advantages of BLAUD'S PILLS over all other ferruginous preparations, and I regard them as the best anti-chlorotic."—Dr. DOUBLET, Ex-President of the Academy of Medicine.

"Of all the ferruginous preparations, which have given good results in the treatment of chlorotic affections, BLAUD'S PILLS appear to us to deserve to hold the first rank."—Dictionnaire universel de médecine, Vol. II, p. 99.

These pills, prepared according to the genuine formulae of the originator, by his nephew, Aug. Blaud, Pharmacist of the Faculty of Paris, are sold only in bottles of 200 pills and half bottles of 100 pills, at a cost of 6 and 3 francs (\$1.00, and seventy-five cents), and never in smaller quantities. See that his name is stamped on each pill. PARIS, 8 RUE PAYENNE, and at every Pharmacy. (Beware of imitations.)

**MEDICO-CHIRURGICAL COLLEGE
OF PHILADELPHIA.**

WINTER SESSION will begin Wednesday, October 1st, and continue until April 16th. PRELIMINARY SESSION begins September 8th. SPRING TERM, April 20th, 1891.

The curriculum is graded, and a preliminary examination and three Annual Winter Sessions are required. Laboratory instruction in Chemistry, Histology, Pathology, Hygiene, Physiology, with Bedside instruction in Medicine, Surgery and Gynecology, is a part of the regular course.

Fees: Matriculation \$5.00. First and second years, each \$75.00. Third year \$100.00. Fourth year free to those in attendance three sessions; to all others \$100.00.

For Announcement or information apply to

**E. E. MONTGOMERY, M.D.,
1818 Arch Street, Philadelphia, Pa.**

**THE JEFFERSON MEDICAL COLLEGE
OF PHILADELPHIA.**

The 68th Annual Session of the Jefferson Medical College begins October 1st and continues nearly 7 months. Preliminary Lectures will be held from 22d of September.

Each student is immediately and personally taught in Obstetrics and Gynecology, Physical Diagnosis, Laryngology, Ophthalmology, Medicinal Chemistry, Pharmacy, Materia Medica, and Experimental Therapeutics; Histology and Experimental Physiology, Minor Surgery, Bandaging, Operations on the Cadaver, Pathology, Neurology, and Electro-Therapeutics.

Three annual regular sessions are required. Bedside instruction in Medicine, Gynecology, Surgery and Obstetrics is a part of the third year course. No extra fee.

The Annual Announcement will be sent on application to **J. W. HOLLAND, M. D., Dean.**

SCOTT'S EMULSION

VERSUS

PLAIN COD LIVER OIL.

Plain Cod Liver Oil is indigestible, deranges the stomach, destroys the appetite, is not assimilated, and in a majority of cases is detrimental to the patient.

SCOTT'S EMULSION

Can be digested in nearly all cases, is assimilated, does not derange the stomach, nor overtax the digestive functions, and can be taken for an indefinite period when the plain cod liver oil cannot be tolerated at all, and with most marked results in *Anæmia, Consumption* and all wasting conditions. It also contains the Hypophosphites of Lime and Soda with Glycerine, which are most desirable adjuncts.

WHEN PHYSICIANS TRY IT THEY INVARIABLY USE IT

in preference to the plain cod liver oil or other *so-called Emulsions* that invariably separate, and hence their integrity and value is destroyed. *Scott's Emulsion is palatable and absolutely permanent* hence its integrity is always preserved.

The formula for Scott's EMULSION is 50 per cent. of the finest Norwegian Cod Liver Oil, 6 grains Hypophosphite of Lime and 3 grains Hypophosphite of Soda to the fluid ounce, Emulsified, or digested to the condition of assimilation with chemically pure Glycerine and Mucilage.

We also wish to call your attention to the following preparations:

CHERRY MALT PHOSPHITES.

A combination of the tonic principles of *Prunus Virginiana*, Malted Barley, Hypophosphites of Lime and Soda, and Fruit Juices. An elegant and efficient brain and nerve tonic.

BUCKTHORN CORDIAL (Rhamnus Frangula).

Prepared from carefully selected German Buckthorn Bark, Juglans Bark and Aromatics. The undoubted remedy for Habitual Constipation.

Be sure and send for samples of the above—delivered free.

SCOTT & BOWNE, 132 South Fifth Avenue, NEW YORK.

THE ATMOSPHERIC TRACTOR.

THE MOST BENEFICENT INVENTION OF THE CENTURY.

IT ENABLES parturition to be rendered almost PAINLESS, and REDUCES the expulsive stage to minutes, instead of hours.

Powerful for good, and powerless to inflict injury on either the mother or the child.

**LIGHT, DURABLE AND ASEPTIC.
INDISPENSABLE TO EVERY PHYSICIAN.**

PRICE, \$3.00.

Sent free by mail or express on receipt of \$3.00. Address

The Atmospheric Tractor Co.,

607 CHESTNUT STREET,

PHILADELPHIA.

CELERINA

NERVE TONIC, STIMULANT AND ANTISPASMODIC.

FORMULA.—Every Fluid Drachm represents FIVE grains EACH—Celery, Coca, Kola, Viburnum and Aromatics.

INDICATIONS.—Loss of Nerve Power (so usual with Lawyers, Preachers, Writers and Business Men), Impotency, Spermatorrhea, Nervous Headache, Neuralgia, Paralysis, Hysteria, Opium Habit, Inebriety, Dyspepsia, and ALL LANCUID conditions of the System.

Indispensable to restore a patient after alcoholic excess.

DOSE.—One or two Teaspoonfuls three or more times a day, as directed by the Physician.

ALETRIS CORDIAL

UTERINE TONIC AND RESTORATIVE.

PREPARED FROM THE ALETRIS FARINOSA OR TRUE UNICORN.

INDICATIONS.—Amenorrhea, Dysmenorrhea, Leucorrhea, Prolapsus Uteri, Sterility, to PREVENT Miscarriage, Etc.

DOSE.—One Teaspoonful three or four times a day.

Unrivaled as a Uterine Tonic in Irregular, Painful, Suppressed and Excessive Menstruation. It Restores Normal Action to the Uterus, and Imparts Vigor to the Entire Uterine System.

Where Women have miscarried during previous pregnancies, or in any case where miscarriage is feared, ALETRIS CORDIAL is Indicated, and should be continuously administered during entire gestation.

S. H. KENNEDY'S

CONCENTRATED EXTRACT OF

PINUS CANADENSIS

DARK.

A NON-ALCOHOLIC LIQUID.

WHITE.

A MOST VALUABLE NON-IRRITATING MUCOUS ASTRINGENT.

INDICATIONS.—Albuminuria, Diarrhea, Dysentery, Night-Sweats, Hemorrhages, Profuse Expectoration, Catarrh, Sore Throat, Leucorrhea, and other Vaginal Diseases, Piles, Sores, Ulcers, Burns, Scalds, Conorrhcea, Cleet, Etc.

When Used as an Injection, to Avoid Staining of Linen, the WHITE Pinus should be used,

RECOMMENDED BY PROMINENT EUROPEAN AND AMERICAN PHYSICIANS.

The above preparations are prepared exclusively for Physicians' Prescriptions, and a sample of each or all of them will be sent to any Physician who wishes to test them, if he will pay the Express Charges.

RIO CHEMICAL CO., St. Louis, Mo., U. S. A.

LONDON.

PARIS.

CALCUTTA.

MONTREAL.

**LIQUOR SANTAL FLAV.
CUM
BUCHU ET CUBEBA.**
(ORIGINAL FORMULA.)

This elegant preparation has been very successfully used in the treatment of long-standing Gonorrhœa and chronic Gleet.

If given in the early stages of Gonorrhœa, it will usually arrest the progress of the disease.

It is especially effective in all inflammations of genito-urinary mucous membranes.

It is perfectly miscible in water. Dose, $\frac{f}{3}$ i to $\frac{f}{3}$ ij, three times a day in water or milk. When prescribing, please specify Liq. Santal Flav. cum Buchu et Cubeba (Ogden's).

**JOHN OGDEN,
CHEMIST,
WALNUT & THIRTEENTH STREETS,
PHILADELPHIA.**

Every Physician in THE WORLD should Read This Advertisement.

**THE RIP VAN WINKLE RECLINING ROCKER.
OFFICE AND GYNAECOLOGICAL CHAIR COMBINED.**

Special Inducements and Extra Discount Given to Physicians.


No. 2.
It is the most wonderful Chair in the world. You can sit in the Chair with your feet on the foot-rest and rock yourself, and it is impossible for a beholder to tell what makes the Chair rock. It is the finest invalid's Chair in the world.


No. 2.
As an Invalid's Chair, a person can recline the back by quarter-inches until he lies straight or horizontally, and without taking his weight from the back of the chair in the least, and CAN ROCK AS WELL LYING DOWN AS SITTING UP. Upholstered in Plush, Leather or Wilton Rugs. This Chair has over 100 changes of position, and sells for \$25.00.

This Chair can be converted from a splendid Office Chair (see cuts Nos. 1 and 3) into one of the most complete Gynaecological Chairs in a half minute, (see cuts 4, 4, and 5).

As a Gynaecological Chair it will take all the positions of the high-priced chairs costing two and three times what this one does.

The Rip Van Winkle Chair has over 100 changes of position. Every chair fully warranted.



No. 3.

The Chair makes 5 pieces of Furniture.

I have attachments by which it can be converted into an Invalid's Rolling Chair in one minute. Parties can get these attachments from factory at any time. Send for full particulars, price-list, and illustrated catalogue showing the principal positions and charges. The Catalogue contains the "Legend of Rip Van Winkle."

Without the Gynaecological Attachment these Chairs

This Chair is sold with or without the Gynaecological attachment, as desired. It is unquestionably the most complete and comfortable Reclining or Lounging Chair ever made. I make a special price to physicians, and also give them a handsome commission on all orders I receive through them for Chairs for their patients. No Chair in the world is so well adapted to ladies suffering from female weakness, and the commission I give on your orders will soon pay for your own Chair.



No. 5.

Make an Elegant Wedding or Holiday Present.
Manufactured by P. C. LEWIS, Box 154, Catskill, N. Y.

WESTERN PENNSYLVANIA MEDICAL COLLEGE
CITY OF PITTSBURGH.

SESSIONS OF 1890-91.

The REGULAR SESSION begins on the last Tuesday of September, and continues six months. During this session, in addition to four Didactic Lectures, two or three hours are daily allotted to Clinical Instruction. Attendance upon two regular courses of lectures is requisite for graduation. A three years' graded course is also provided. The SPRING SESSION embraces recitations, clinical lectures and exercises, and didactic lectures on special subjects; this session begins the second Tuesday in April, and continues ten weeks.

The laboratories are open during the collegiate year for instruction in chemistry, microscopy, practical demonstrations in medical and surgical pathology, and lessons in normal histology. Special importance attaches to "the superior clinical advantages possessed by this College." For particulars, see annual announcement and catalogue, for which, address the Secretary of Faculty, Prof. T. M. T. McKENNA.

Business correspondence should be addressed to
Prof. W. J. ASDAHL, 3107 Penn Avenue, Pittsburgh.

To Rent.

Apply

1816 Christian Street,
Philadelphia.

For Sale.

Address

Medical and Surgical Reporter,
P. O. Box 843, Philadelphia.

PRACTICE AND
OFFICE of a recently-deceased physician.
Terms low.

A TOEPLER-HOLTZ
STATIC BATTERY, in
case, as good as new. 16
inch revolving plate.

WALNUT LODGE HOSPITAL,
HARTFORD, CONN.

Organized in 1880 for the special medical treatment of
ALCOHOL AND OPIUM INEBRIATES.

Elegantly situated in the suburbs of the city, with every appointment and appliance for the treatment of this class of cases, including Turkish, Russian, Roman, Saline and Medicated Baths. Each case comes under the direct personal care of the physician. Experience shows that a large per cent. of these cases are curable, and all are more or less benefited from the application of exact hygienic and scientific measures. This institution is founded on the well-recognized fact that INEBRIETY is a DISEASE, and CURABLE, and all these cases require rest, change of thought and living, in the best surroundings, together with every means known to science and experience to bring about this result. Applications and all inquiries should be addressed

T. D. CROTHERS, M.D.,
Sup't Walnut Lodge, Hartford, Conn.

**DR. MASSEY'S
PRIVATE SANITARIUM**

212 South Fifteenth Street

PHILADELPHIA

This institution, in addition to complete arrangements for the treatment of Nervous Diseases by rest, electricity, massage, etc., under comfortable surroundings, is specially equipped for the treatment of cases of Fibroid Tumors and other intractable diseases of the pelvic viscera, by the conservative use of strong electric currents. For particulars, address

DR. G. BETTON MASSEY,
Physician in Charge.

Mellin's Food FOR INFANTS AND INVALIDS.

A SOLUBLE DRY EXTRACT, prepared from Malted Barley and Wheat, consisting of Dextrin, Maltose, Albuminates, and Salts.

The SUGAR in MELLIN'S FOOD is MALTOSE. MALTOSE is the PROPER SUGAR for use in connection with cow's milk.

The sugar formed by the action of the Ptyalin of the Saliva and the Amylopsin of the Pancreas upon starch is MALTOSE. In the digestive tract MALTOSE is absorbed UNCHANGED.

—Landois and Sterling.

MALTOSE is a saccharose, not a glucose, and is a form of sugar which does not ferment.

—Materia Medica and Therapeutics, Dr. Mitchell Bruce.

"I have never seen any signs of fermentation which I could attribute to the influence of MALTOSE."

—Eustace Smith, M.D., F.R.C.S.

MELLIN'S FOOD, prepared according to the directions, is a true LIEBIG'S FOOD and the BEST SUBSTITUTE for Mother's Milk yet produced.

IT REQUIRES NO COOKING.

THE DOLIBER-GOODALE CO.,

BOSTON, MASS.

PHILADELPHIA POLYCLINIC AND COLLEGE FOR GRADUATES IN MEDICINE.

THE POLYCLINIC HOSPITAL,

NORTHWEST CORNER BROAD AND LOMBARD STS.
REGULAR SESSIONS RECOMMENCE SEPT. 15TH

Pupils may begin at any time. Tickets good for Six Weeks' Daily Clinics from date of issue or One Clinic Weekly, for three months.

PRACTICAL, INDIVIDUAL INSTRUCTION, TO PHYSICIANS ONLY, IN ALL BRANCHES OF MEDICINE AND SURGERY.

FOR ANNOUNCEMENT, ADDRESS,
SOLOMON SOLIS-COHEN, M.D., SECRETARY.

NEW YORK POST-GRADUATE MEDICAL SCHOOL AND HOSPITAL.

SESSIONS 1890-91.

The Autumn and Winter Sessions of the Post-Graduate Medical School opened on September 15. The Hospital is fully utilized for clinical purposes. There are ninety beds, including thirty for babies in the adjacent building. No Post-Graduate Medical School in this country can offer such opportunities for seeing important surgical and gynecological operations under its own roof as does this institution. Besides, regular clinics, medical and surgical, are held by professors and instructors in those general and special hospitals with which they are connected, which comprises nearly all the great hospitals of the city. The dispensary of the school and other large dispensaries furnish ample means of instruction in every class of walking cases.

Instruction is given in every branch of medicine and surgery. The laboratory is large and well equipped. Surgical anatomy and operative surgery are thoroughly taught. The instruction is personal, the practitioner being brought in direct contact with the patient. No other courses in any department will be required if those to be had in the Post-Graduate School are taken. The faculty were the first in the United States to organize private courses for practitioners of medicine, and they pledge themselves to leave nothing undone to make graduates in medicine profit by the unrivaled clinical advantages of the City of New York.

For catalogue and schedule, address

CLARENCE C. RICE, M. D., Secretary.

D. B. St. JOHN ROOSA, M. D., LL.D., PRESIDENT. 226 East 20th St., New York City.
F. E. FARRELL, SUPERINTENDENT.

COLLEGE OF MEDICINE. SYRACUSE UNIVERSITY.

FACULTY.

JOHN VAN DUVN, M. D., Surgery.	GAYLORD P. CLARK, M. D., Anatomy.
HENRY D. DIDAMA, M. D., Practice and Clinical Medicine.	WILLIAM H. DUNLAP, M. D., Dermatology.
WILLIAM T. PLANT, M. D., Diseases of Children.	JOHN L. HEFFRON, M. D., Materia Medica and Therapeutics.
ALFRED MERCER, M. D., State Medicine.	HENRY B. ALLEN, M. D., Obstetrics.
WILLIAM MANLIUS SMITH, M. D., Chemistry.	A. CLIFFORD MERCER, M. D., Pathology.
DAVID M. TOTMAN, M. D., Clinical Surgery.	HENRY L. ELSNER, M. D., Clinical Medicine.
NATHAN JACOBSON, M. D., Clinical Surgery and Laryngology.	FRANK W. MARLOW, M. D., Ophthalmology.

Three Years' Graded Course. Year begins October 7th, 1890, ends in June, 1891.
FIRST YEAR: Anatomy, Physiology, Chemistry, Histology, Materia Medica.

SECOND YEAR: Anatomy, Physiology, Chemistry, Therapeutics, Pathology, Practice, Surgery, Clinical Medicine and Surgery.

THIRD YEAR: Therapeutics, Practice, Surgery, Children, Clinics, Obstetrics, Gynecology, Forensic Medicine, Ophthalmology, Hygiene.

An experience of seventeen years has demonstrated that a three years' graded course is superior in every respect to an ungraded, unnatural, and cramming course of two winters.

Tuition, \$80.00 for each Year. Perpetual Ticket, \$200.00. Diploma, \$25.00.

For circulars with full information, address

W. H. DUNLAP, M. D., Registrar, SYRACUSE, N. Y.

UNIVERSITY OF PENNSYLVANIA.—Medical Department.

The 125TH ANNUAL WINTER SESSION will begin Wednesday, October 1st, 1890, at 12 M., and will continue seven months. The PRELIMINARY SESSION begins September 22d, 1890, the Spring Term early in May, 1891.

The curriculum is graded and three annual winter sessions are required. Practical instruction, including laboratory work in Chemistry, Histology, Osteology and Pathology, with Bedside Instruction in Medicine, Surgery and Gynecology are a part of the regular course and without additional expense.

FACULTY.


JOSEPH LEIDY, M.D., LL.D., Professor of Anatomy.	WILLIAM F. NORRIS, M.D., Honorary Prof. of Ophthalmology.
D. HAYES AGNEW, M.D., LL.D., Honorary Professor of Clinical Surgery.	BARTON COOK HIRST, M.D., Professor of Obstetrics.
WILLIAM PEPPER, M.D., LL.D., Professor of Theory and Practice of Medicine, and of Clinical Medicine.	J. WILLIAM WHITE, M.D., Professor of Clinical Surgery.
WILLIAM GOODELL, M.D., Professor of Gynecology.	JOHN GUITÉRAS, M.D., Professor of General Pathology and Morbid Anatomy.
JAMES TYSON, M.D., Professor of Clinical Medicine.	GEORGE A. PIERSON, M.D., Professor of Histology and Embryology.
MORATIUS C. WOOD, M.D., LL.D., Professor of Materia Medica, Pharmacy and General Therapeutics.	SAMUEL G. DIXON, M.D., Professor of Hygiene.
THEODORE G. WORMLEY, M.D., LL.D., Professor of Chemistry and Toxicology.	For Catalogue and announcement containing particulars apply to
JOHN ASHURST, JR., M.D., Professor of Surgery and of Clinical Surgery.	DR. JAMES TYSON, Dean, 36th and Woodland Avenue, Philadelphia
EDWARD T. REICHERT, M.D., Professor of Physiology.	

A. G. SPALDING & BROS.,

MANUFACTURERS OF

Gymnasium Goods and Apparatus

OF EVERY DESCRIPTION.



INDIAN CLUBS,
DUMB BELLS,
ROWING MACHINES,
CHEST EXPANDERS,
HORIZONTAL AND
PARALLEL BARS,
STRIKING BAGS,
PULLEY WEIGHTS,
BOXING GLOVES,
FENCING FOILS, Etc., Etc.

The O. K. Peerless Pulley Weight and Machine, combining all the necessary exercise for the perfect development of the chest and body, the most simple and complete method for home exercise and training ever invented.

Uniforms and Clothing for all gymnasium uses. We are the *sole* agents for the only genuine *Shaker Sweater*, a garment indispensable to the *athlete* and in the *gymnasium*. This *Sweater* has but to be seen to be commended as the most superior garment for this purpose in existence.

Use only the *Spalding* line of *Athletic* and *Sporting Goods*, the standard and best. Prices always reasonable.

CATALOGUES FREE.

A. G. SPALDING & BROS.,

CHICAGO,
108 MADISON STREET.

NEW YORK,
241-243 BROADWAY.

PHILADELPHIA,
1022 MARKET STREET.

GRIFFITH & CO.'S Compound Mixture of Guaiac, Stillingia, etc.

For Acute and Chronic Rheumatism, Gout, Lumbago, Neuralgia, and Kindred Complaints.

TO PHYSICIANS.—Gentlemen: We would respectfully draw your attention to our COMPOUND MIXTURE OF GUIAC, STILLINGIA, ETC.

THIS is not a new preparation, but has been in constant use by many prominent practitioners of medicine for several years, and its beneficial results in the treatment of the diseases indicated, including Syphilitic troubles, have been fully established.

When ordering this Preparation, in order to avoid delay or misunderstanding, Physicians will please specify "GRIFFITH & CO.'S," or Physicians in the city can send their patients direct to our pharmacies, at No. 67 Third Avenue, cor. 11th St., or 2241 Third Avenue, cor. 122d St., New York, where, at any time, further information will be cheerfully furnished. Out of town Physicians can order through their druggists or direct from us.

WE have hundreds of testimonials from prominent physicians who have prescribed and personally used this mixture. It is manufactured for **PHYSICIANS' PRESCRIPTIONS** only. Always specify **GRIFFITH & CO.'S**. If you have an obstinate case of Rheumatism under treatment, inclose One Dollar and receive, by express, a regular size bottle, or we will send, upon request, a sample bottle providing you will pay express charges. Wholesale Price-List—8-ounce size, \$10.50 per dozen ; 16-ounce size, \$20.00 per dozen. In lots of one dozen and upward, we prepay express charges to any point east of the Rocky Mountains. (Do not overlook this offer, for you may be pleased and possibly surprised at the results.)

P. S.—The advertising of this article is confined strictly to Medical Journals, and the general verdict of physicians is that if this mixture does not act on a case, it is a difficult matter to find something that will.

Very respectfully,

GRIFFITH & CO., Chemists and Pharmacists,
67 Third Ave., cor. 11th St., and 2241 Third Ave., cor. 122d St., **NEW YORK.**

Carried in stock by the principal Wholesale Druggists in the U. S.

SVAPNIA OR PURIFIED OPIUM

FOR PHYSICIANS USE ONLY.
Contains the Anodyne and Soothing Alkaloids, Codeia, Narceia and Morphia. Excludes the Poisonous and Convulsive Alkaloids, Thebaine, Narcotine and Papaverine.

SVAPNIA has been in steadily increasing use for over twenty years, and whenever used has given great satisfaction.

To PHYSICIANS of REPUTATION, not already acquainted with its merits, samples will be mailed on application.

SVAPNIA is made to conform to a uniform standard of Opium of Ten per cent. Morphia strength.

JOHN PARR, Manufacturing Chemist, New York.
C. J. CRITTENDEN, Cor'l Agnd, 115 Fulton St., N. Y.

To whom all orders for samples must be addressed.

SVAPNIA IS FOR SALE BY DRUGGISTS GENERALLY.

St. Vincent Home Maternity Hospital,

70th Street and Woodland Avenue,

(PASCHALL STATION) PHILADELPHIA.

Conducted by the Sisters of Charity.

INFANTS and Children under two years of age received.

Patients for MATERNITY DEPARTMENT accommodated according to their means.

IN THE WARDS terms range from \$5.00 to \$8.00 per week.

IN THE PRIVATE ROOMS terms range from \$10.00 to \$20.00 per week.

For further information apply to or address

THE SISTER IN CHARGE,

PASCHALVILLE,

PHILADELPHIA, PA.

GOUDRON DE BLOUNT

PREPARED FROM THE GENUINE CAROLINA TAR.

DOSE.—One fluid drachm four or more times a day (as indicated), either full strength, diluted, or, in combination.

INDICATIONS.—Chronic and acute affections of the Air Passages, Coughs, Colds, Bronchitis, Asthma and Consumption.

WILLIAM MURRELL, M.D., F.R.C.P.,

Lecturer on Pharmacology and Therapeutics at the Westminster Hospital; Examiner in Materic Medicæ to the Royal College of Physicians of London; Fellow of the Medical-Chirurgical College of Philadelphia.
Says:—"I have used with success 'Goudron de Blount.' The results have been good, and the preparation is popular with patients."

~~PREPARED EXCLUSIVELY FOR PHYSICIANS' PRESCRIPTIONS BY~~

R. E. BLOUNT, 33 RUE ST. ROCH, PARIS.

WHOLESALE AGENTS FOR UNITED STATES AND CANADA,

BATTLE & CO.,

CHEMISTS' CORPORATION,

Clinical Notes sent on Application.

ST. LOUIS, MO.

FAIRCHILD'S ESSENCE OF PEPSINE

Contains the essential active principle of the gastric juice,
obtained by direct extraction from the rennet.

It is by far the most active, reliable, and agreeable of all the preparations of pepsin.

This preparation has been ten years before the medical profession, and enjoys a high reputation and a very wide use.

There can be no doubt that the value of the digestive ferment is enhanced by skilful and judicious combination with aromatics, and this Essence of Pepsine is peculiarly grateful and stomachic.

It tends to promote the natural digestive functions, and has been found in practice to be especially suited to digestive disorders of infants, for the correction of flatulency, indigestion, etc.

It also finds practical use in the preparation of junket as an agreeable food for invalids, and in the preparation of whey as a temporary food for infants.

We respectfully request that physicians who prefer Fairchild's preparation will kindly specify them in order to avoid the imitations which are now urged upon the market as "just as good as Fairchild's." The substitution of these imitations can certainly be of no advantage whatever except to the maker of them and the one who foists them upon the consumer, as the same price is charged for these imitations as for the genuine article.

FAIRCHILD BRO'S & FOSTER,

82 and 84 Fulton Street. New York.

"What a boon it would be to the Medical Profession if some reliable Chemist would bring out an Extract of Malt in combination with a well-digested or peptonized Beef, giving us the elements of Beef and the stimulating and nutritious portions of Ale."—J. MILNER FOTHERGILL, M.D.

Ale & Beef

"PEPTONIZED"

(EXTRACT: BOVIS GUM MALTO.)

Is the identical combination suggested by the late eminent Fothergill.

**EACH BOTTLE REPRESENTS 1-4 POUND
OF LEAN BEEF THOROUGHLY PEPTONIZED.**

It is the only Mild Stimulant combined with a Perfect Food known. It is very palatable, aids digestion, is retained by the most delicate stomach, and the purity of its ingredients is guaranteed.

THE "PEPTONIZED" BEEF

is manufactured by Prof. Preston B. Rose, of Chicago, late of the faculty of the University of Michigan.

THE ALE USED IS THE PUREST AND BEST MADE IN AMERICA

Especially brewed for this purpose, and guaranteed to be equal to the best imported ales, as only the best Canada Malt and the choicest new hops are used in its manufacture.

IT IS A REAL FOOD; NOT A MERE STIMULANT,

as it contains all the albumen and fibrin of the beef as well as the nutritive qualities of the malted barley.

It is most useful during the period of Gestation, in allaying all vomiting, and invaluable to Nursing Mothers; also in Typhoid Fever and Dysentery.


Prof. G. A. Liebig says: "A careful chemical examination of the Peptonized Ale and Beef shows a much larger per cent. of nitrogenous blood and muscle-making matter over all other malt extracts, and that it is also rich in Diastase, giving it the power to digest Starch Foods."

PREPARED BY

The Ale and Beef Company,

DAYTON, OHIO, U.S.A.

Two full-sized bottles will be sent FREE to any physician who will pay express charges.
Please mention this journal.





MEDICAL AND SURGICAL REPORTER

A WEEKLY JOURNAL.

ESTABLISHED IN 1853 BY S. W. BUTLER, M.D.

CHARLES W. DULLES, M. D., EDITOR AND PUBLISHER,

Entered as Second-Class matter at Philadelphia P. O.

N. E. COR. 18TH & WALNUT STS., PHILADELPHIA

Vol. LXIII, No. 26.
Whole No. 1765.

DECEMBER 27, 1890.

\$5.00 per Annum.
10 Cents a Copy.

CONTENTS:

CLINICAL LECTURES.

- GOODELL, WILLIAM, M. D., Philadelphia, Pa.—
Closure of Vulva for Vesico-Vaginal Fistula.—
Fibroid Tumors..... 728
MUSSER, JOHN H., M. D., Philadelphia, Pa.—Hydatid Cyst of the Liver.—Pleurisy Associated with Tuberculosis..... 726

COMMUNICATIONS.

- ATLEE, JOHN E., M. D., Chattanooga, Tenn.—Case of Ovariotomy..... 728
VINTON, MARIA MITCHELL, A. M., M. D., New York.—Carcinoma of the Gall-bladder and Liver from Biliary Calculi..... 729

PERISCOPE.

- Diabetic Coma.—Salinaphthol in Rheumatism.—Railway Readers.—Ergotin in Gleet..... 731-732

EDITORIALS.

- THE END OF THE YEAR..... 733
HYSTERIA IN MALES..... 733
A KOCH INSTITUTE IN NEW YORK..... 734

BOOK REVIEWS.

- KEATING; Cyclopaedia of the Diseases of Children 734

SPECIAL ARTICLE.

- Koch's Remedy for Tuberculosis..... 735

NOTES AND COMMENTS.

- Exact Dosage in the Cataphoretic Use of Drugs.—Pleasant Laxative for Infants.—Evidences of Criminal Poisoning.—Treatment of Delirium Tremens.—Reduction of Hip Dislocations.—Bouducine as a Febrifuge.—The Quantity of Blood in the Brain.—Hydrazole Acid..... 738-740

FOR THE CURE OF NERVOUS HEADACHES

SEDATIVE EFFERVESCENT

EFFERVESCENT ANODYNE




R Caffein 1 grain, Brom. Soda 30 grains in each heaping teaspoonful.

Useful in Nervous Headache, Sleeplessness, Excessive Study, Migraine, Nervous Debility, Mania, as a Remedy in Seasickness and Epilepsy.

Dose and Composition.—A heaping teaspoonful, containing Bromo-Soda 30 grs., and Caffein 1 gr., in half a glass of water, to be repeated once after an interval of thirty minutes if necessary.

As the composition of the above effervescent salts is such as physicians can use with confidence, and you will please specify **Bromo-Soda** or **Bromo-Potash**, **Warner & Co.**, when ordering or prescribing to avoid the substitution of other "Bromos." Prepared only by



R Caffein 1 grain, Bromide Potash 20 grains in each heaping teaspoonful.

Useful in Nervous Headache, Sleeplessness, Excessive Study, Migraine, Nervous Debility, Mania, as a Remedy in Seasickness and Epilepsy.

Physicians desiring the Potash Salt can obtain the same by ordering or prescribing **Bromo-Potash** (**Warner & Co.**), the composition of which is: Brom. Potash 20 grs., Caffein 1 gr.

WILLIAM R. WARNER & CO.,
1228 MARKET STREET, PHILADELPHIA. 18 LIBERTY STREET, NEW YORK.

JOHN C. BAKER & CO.'S STANDARD PREPARATIONS.

BAKER'S PURE COD LIVER OIL NORWEGIAN.

BAKER'S EMULSION WITH HYPOPHOSPHITES OF LIME AND SODA.

BAKER'S C. L. OIL WITH EXT. OF MALT.

SOL. FERROUS MALATE A. P. M. CO.

SOL. SALICYLATE OF IRON A. P. M. CO.

FER. MALATE, GENTIAN AND CINCHONA A. P. M. CO.

LITERATURE MAILED ON REQUEST.

JOHN C. BAKER & CO., PHILADELPHIA, PA.



ANTISEPTICS, DISINFECTANTS AND OXIDANTS.

"Sanitas" is prepared by Oxidizing Terpene in the presence of Water
with Atmospheric Air.

"SANITAS" DISINFECTING FLUID.

An aqueous extract of Air-Oxidized Terpene. Its active principles include Soluble Camphor ($C_{10}H_{16}O_3$), Peroxide of Hydrogen, and Thymol. Invaluable to the physician for internal or external application.

"SANITAS" DISINFECTING OIL.

Air-Oxidized Terpene. Its active principle is Camphoric Peroxide ($C_{10}H_{16}O_3$), a substance which produces Peroxide of Hydrogen when placed in contact with water or moist surfaces (wounds, mucous membranes, and other tissues). It is a powerful germicidal and oxidizing agent. For fumigations and inhalations in the treatment of diseases of the respiratory organs the oil only requires to be evaporated from boiling water.

"Sanitas" is Fragrant, Non-poisonous, and does not stain or corrode. It is put up in the form of FLUIDS, OIL, POWDERS AND SOAPS.

For Reports by Medical and Chemical Experts, Samples, Prices, etc., apply to the Factory,

636, 638, 640, and 642 West 55th Street, New York.

SIR MORELL MACKENZIE, in the *Journal of Laryngology*, published in London, says the following:

"Most of the German Saline Springs are situated on a range of the Taunus Hills. Of these, unquestionably one of *the very best* is

SODEN.

"Containing, as it does, some 23 springs, with a temperature varying between 12° to 18° C., the waters of which are charged with Chloride of Sodium in varying degrees up to 116 Grammes per litre, and some of which are further charged with Carbonic Acid, the Health Resort of Soden is unquestionably one of the very best places for the treatment of Pulmonary and Catarrhal complaints.

"COMPRESSED PASTILLES of the Soden Water are now an article of commerce, as well as the mineral waters. These are made from two of the well-waters, and containing a large amount of Chloride of Sodium, are particularly serviceable in pharyngeal catarrhs, and may even in some degree be used where it is desired to obtain the effect of the Soden treatment, in persons who are unable to make the necessary journey to the Spa itself. I have watched the effects of the Soden Waters for a considerable period, and regard them as extremely valuable in obstinate catarrhal affections of the throat. The small amount of iron which they contain renders them very useful in the stages of Throat Consumption, and they do good in nearly all cases of relaxation of the mucous membrane.

"The Pastilles offer a most convenient method of using the Waters; producing both a local and general effect. They are especially beneficial in catarrhal diseases of the air-passages; I have frequently found them of great service in the case of singers and public speakers.

(Signed)

"MORELL MACKENZIE, M. D., LONDON."

DR. NORRIS WOLFENDEN, Senior Physician Throat Hospital, London, writes:

"SIR:—The Wells of Soden are known to possess most powerful medicinal virtues, and have been found of the utmost benefit in the treatment of Catarrhal conditions of the upper respiratory tract, such as Bronchitis and Emphysema, and even in Pulmonary Phthisis. Many chronic throat conditions are equally benefited. Possessing a large amount of Sodium Chloride in their composition, they have a character which is quite special, and the fact that the several Wells of Soden contain varying amounts of this important mineral constituent, renders it possible to treat many complaints of opposite nature on the same spot.

"I consider the 'Soden Mineral Pastilles,' which you have introduced to me, to be a most convenient and pleasant mode of obtaining the medicinal benefits of the Soden Wells. They form an agreeable and effective lozenge, and I feel certain will be of great service to patients afflicted with certain forms of chronic throat disease. Sprays of the same waters would, I am sure, be of much service. I hope your efforts to introduce the Soden Mineral Waters and Pastilles into this country will meet with success.

"I am, Sir, yours faithfully,

(Signed) "R. NORRIS WOLFENDEN, M. D. Cantab.,
"Senior Physician Throat Hospital, London; Editor of *Journal of Laryngology*."

SAMPLES AND PAMPHLETS MAILED UPON APPLICATION.

Soden Mineral Springs Co., Ltd., Sole Agents,
P. O. BOX 3319, NEW YORK.

Doctor:

You have often noticed that many patients, after a course of typhoid fever, erysipelas, diphtheria, and kindred diseases, undergo a long, tedious convalescence. For many weary weeks they sit around the house or drag their emaciated bodies feebly along. Their flame of life is so dim that it is easily puffed out by any slight indisposition or intercurrent malady, that may chance to attack them, and hence we see so many *cured* cases become easy victims by the predisposition which their debility constitutes.

Now, what is the true condition in such as case as this?

Why, simply that in the febrile process immense hosts of the tissue cells have been consumed or carried away, and most of those remaining are worn out and only await a vigorous circulation to remove them and deposit healthy cells in their places.

In addition to the emaciation you notice that the tissues still left are flabby and apparently lifeless. The nutritive organs are so weakened that they are barely sufficient to maintain life without furnishing the required new material to carry on the process of repair. Thus the individual remains an easy prey to pathogenic influences which one in robust health can successfully resist.

Can anything be done for such cases?

Certainly.

But you would better leave them to the unassisted reparative powers of nature than to dose them with the irritant alleged "tonics" now so much in vogue. The organs of nutrition only want to be supplied with true cell and tissue food in an assimilable form.

This you will find in McArthur's Compound Syrup of the Hypophosphites of Lime and Soda. This is a CHEMICALLY PURE preparation, a quality so rare in Hypophosphite preparations. It is prepared in such a manner that the important ingredients go entirely to the rapid rebuilding of the tissues.

Give this to your convalescent patient and you will see a remarkable increase in flesh, strength, and buoyancy of spirits. The convalescence will be greatly shortened, and in every way the Hypophosphites will prove to them to be a true Elixir of Life.

You can also make an enviable reputation for yourself by prescribing it in Phthisis and all Tubercular affections, Scrofula, Throat, and Bronchial diseases, Nervous disorders and degenerations, and Fibroid Tumors.

Send for our pamphlet on "The Curability and Treatment of Consumption" (free), and a full size bottle of McArthur's Syrup, free, if you will agree to pay express charges.

McARTHUR HYPOPHOSPHITE CO.,

BOSTON, MASS.

376
2

